

ACOT8 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP50587**Specification**

ACOT8 Antibody - Product Information

Application	IF, WB
Primary Accession	O14734
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36 KDa
Antigen Region	163-189

ACOT8 Antibody - Additional Information**Gene ID** 10005**Other Names**

Acyl-coenzyme A thioesterase 8, Acyl-CoA thioesterase 8, Choloyl-coenzyme A thioesterase, HIV-Nef-associated acyl-CoA thioesterase, PTE-2, Peroxisomal acyl-coenzyme A thioester hydrolase 1, PTE-1, Peroxisomal long-chain acyl-CoA thioesterase 1, Thioesterase II, hACTE-III, hACTEIII, hTE, ACOT8, ACTEIII, PTE1, PTE2

Dilution

IF~~1:100
WB~~ 1:1000

Format

Rabbit IgG in phosphate buffered saline (without Mg²⁺ and Ca²⁺), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

Storage Conditions

-20°C

ACOT8 Antibody - Protein Information**Name** ACOT8**Synonyms** ACTEIII, PTE1 {ECO:0000303|PubMed:100925}**Function**

Catalyzes the hydrolysis of acyl-CoAs into free fatty acids and coenzyme A (CoASH), regulating their respective intracellular levels (PubMed:15194431, PubMed:9153233, PubMed:9299485). Displays no strong substrate specificity with respect to the carboxylic acid moiety of Acyl-CoAs (By similarity). Hydrolyzes medium length (C2 to C20)

straight-chain, saturated and unsaturated acyl-CoAS but is inactive towards substrates with longer aliphatic chains (PubMed:9153233, PubMed:9299485). Moreover, it catalyzes the hydrolysis of CoA esters of bile acids, such as choloyl-CoA and chenodeoxycholoyl-CoA and competes with bile acid CoA:amino acid N-acyltransferase (BAAT) (By similarity). Is also able to hydrolyze CoA esters of dicarboxylic acids (By similarity). It is involved in the metabolic regulation of peroxisome proliferation (PubMed:15194431).

Cellular Location

Peroxisome matrix. Note=Predominantly localized in the peroxisome but a localization to the cytosol cannot be excluded

Tissue Location


Detected in a T-cell line (at protein level). Ubiquitous (PubMed:9153233, PubMed:9299485)

ACOT8 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ACOT8 Antibody - Images



Immunofluorescence analysis of A549 cells, using ACOT8 antibody.



Western blot analysis of lysate from Raji cell line, using ACOT8 Antibody (AP50587). AP50587 was diluted at 1:1000. A goat anti-rabbit IgG H&L (HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.

ACOT8 Antibody - Background

Acyl-CoA thioesterases are a group of enzymes that catalyze the hydrolysis of acyl-CoAs to the free fatty acid and coenzyme A (CoASH), providing the potential to regulate intracellular levels of acyl-CoAs, free fatty acids and CoASH. May mediate Nef-induced down-regulation of CD4. Major thioesterase in peroxisomes. Competes with BAAT (Bile acid CoA: amino acid N- acyltransferase) for bile acid-CoA substrate (such as chenodeoxycholoyl-CoA). Shows a preference for medium-length fatty acyl-CoAs (By similarity). May be involved in the metabolic regulation of peroxisome proliferation.

ACOT8 Antibody - References

- Watanabe H., et al. *Biochem. Biophys. Res. Commun.* 238:234-239(1997).
- Liu L.X., et al. *J. Biol. Chem.* 272:13779-13785(1997).
- Jones J.M., et al. *J. Biol. Chem.* 274:9216-9223(1999).
- Deloukas P., et al. *Nature* 414:865-871(2001).
- Ishizuka M., et al. *Exp. Cell Res.* 297:127-141(2004).