

**G6PD Antibody (Center)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP5094c**

**Specification**

---

**G6PD Antibody (Center) - Product Information**

Application	<b>WB, IHC-P, FC,E</b>
Primary Accession	<a href="#">P11413</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Calculated MW	<b>59257</b>
Antigen Region	<b>297-326</b>

**G6PD Antibody (Center) - Additional Information**

**Gene ID** 2539

**Other Names**

Glucose-6-phosphate 1-dehydrogenase, G6PD, G6PD

**Target/Specificity**

This G6PD antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 297-326 amino acids from the Central region of human G6PD.

**Dilution**

WB~~1:1000  
IHC-P~~1:50~100  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

G6PD Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**G6PD Antibody (Center) - Protein Information**

**Name** G6PD

**Function** Catalyzes the rate-limiting step of the oxidative pentose- phosphate pathway, which

represents a route for the dissimilation of carbohydrates besides glycolysis. The main function of this enzyme is to provide reducing power (NADPH) and pentose phosphates for fatty acid and nucleic acid synthesis.

#### Cellular Location

Cytoplasm, cytosol. Membrane; Peripheral membrane protein

#### Tissue Location


Isoform Long is found in lymphoblasts, granulocytes and sperm

### G6PD Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### G6PD Antibody (Center) - Images



Western blot analysis of G6PD Antibody (Center) (Cat. #AP5094c) in K562, MCF-7 cell line lysates (35ug/lane). G6PD (arrow) was detected using the purified Pab.



G6PD Antibody (Center) (Cat. #AP5094c) IHC analysis in formalin fixed and paraffin embedded testis followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the G6PD Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



G6PD Antibody (Center) (Cat. #AP5094c) flow cytometric analysis of MCF-7 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

### **G6PD Antibody (Center) - Background**

G6PD encodes glucose-6-phosphate dehydrogenase. This protein is a cytosolic enzyme encoded by a housekeeping X-linked gene whose main function is to produce NADPH, a key electron donor in the defense against oxidizing agents and in reductive biosynthetic reactions. G6PD is remarkable for its genetic diversity. Many variants of G6PD, mostly produced from missense mutations, have been described with wide ranging levels of enzyme activity and associated clinical symptoms. G6PD deficiency may cause neonatal jaundice, acute hemolysis, or severe chronic non-spherocytic hemolytic anemia.

### **G6PD Antibody (Center) - References**

Louicharoen, C., et al. *Science* 326(5959):1546-1549(2009)  
 Zhong, D.N., et al. *Zhongguo Dang Dai Er Ke Za Zhi* 11(12):970-972(2009)  
 Tiono, A.B., et al. *Am. J. Trop. Med. Hyg.* 81(6):969-978(2009)

### **G6PD Antibody (Center) - Citations**

- [Activation of pro-survival metabolic networks by 1,25\(OH\) does not hamper the sensitivity of breast cancer cells to chemotherapeutics.](#)