

**E Cadherin Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP51064**

**Specification**

---

**E Cadherin Antibody - Product Information**

Application	<b>WB, IHC-P, E</b>
Primary Accession	<a href="#">P12830</a>
Reactivity	<b>Human, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>100 KDa</b>

**E Cadherin Antibody - Additional Information**

**Gene ID** 999

**Other Names**

Cadherin-1, CAM 120/80, Epithelial cadherin, E-cadherin, Uvomorulin, CD324, E-Cad/CTF1, E-Cad/CTF2, E-Cad/CTF3, CDH1, CDHE, UVO

**Format**

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**E Cadherin Antibody - Protein Information**

**Name** CDH1 ([HGNC:1748](#))

**Function**

Cadherins are calcium-dependent cell adhesion proteins (PubMed:<a href="http://www.uniprot.org/citations/11976333" target="\_blank">11976333</a>). They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. CDH1 is involved in mechanisms regulating cell-cell adhesions, mobility and proliferation of epithelial cells (PubMed:<a href="http://www.uniprot.org/citations/11976333" target="\_blank">11976333</a>). Has a potent invasive suppressor role. It is a ligand for integrin alpha-E/beta-7.

**Cellular Location**

Cell junction, adherens junction. Cell membrane; Single-pass type I membrane protein. Endosome. Golgi apparatus, trans-Golgi network. Cytoplasm {ECO:0000250|UniProtKB:P09803}. Cell junction {ECO:0000250|UniProtKB:Q90Z37}. Note=Colocalizes with DLGAP5 at sites of cell-cell contact in intestinal epithelial cells. Anchored to actin microfilaments through association with alpha-, beta- and gamma- catenin. Sequential proteolysis induced by apoptosis or calcium influx, results in translocation from sites of cell-cell contact to the cytoplasm. Colocalizes with RAB11A endosomes during its transport from the Golgi apparatus to the plasma membrane

**Tissue Location**

Expressed in granuloma macrophages (at protein level) (PubMed:27760340). Expressed in the liver (PubMed:3263290)

**E Cadherin Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**E Cadherin Antibody - Images****E Cadherin Antibody - Background**

Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. CDH1 is involved in mechanisms regulating cell-cell adhesions, mobility and proliferation of epithelial cells. Has a potent invasive suppressor role. It is a ligand for integrin alpha-E/beta-7.

**E Cadherin Antibody - References**

Bussemakers M.J.G., et al. Mol. Biol. Rep. 17:123-128(1993).  
Oda T., et al. Proc. Natl. Acad. Sci. U.S.A. 91:1858-1862(1994).  
Rimm D.L., et al. Biochem. Biophys. Res. Commun. 200:1754-1761(1994).  
Ito K., et al. Oncogene 18:7080-7090(1999).  
Ota T., et al. Nat. Genet. 36:40-45(2004).