

gp91 phox Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51139

Specification

gp91 phox Antibody - Product Information

Application	WB, E
Primary Accession	P04839
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	65 KDa

gp91 phox Antibody - Additional Information

Gene ID 1536

Other Names

Cytochrome b-245 heavy chain, 1---, CGD91-phox, Cytochrome b(558) subunit beta, Cytochrome b558 subunit beta, Heme-binding membrane glycoprotein gp91phox, NADPH oxidase 2, Neutrophil cytochrome b 91 kDa polypeptide, Superoxide-generating NADPH oxidase heavy chain subunit, gp91-1, gp91-phox, p22 phagocyte B-cytochrome, CYBB, NOX2

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

gp91 phox Antibody - Protein Information

Name CYBB ([HGNC:2578](#))

Synonyms NOX2

Function

Catalytic subunit of the phagocyte NADPH oxidase complex that mediates the transfer of electrons from cytosolic NADPH to O₂ to produce the superoxide anion (O₂⁻) (PubMed: [15338276](http://www.uniprot.org/citations/15338276), PubMed: [36241643](http://www.uniprot.org/citations/36241643), PubMed: [36413210](http://www.uniprot.org/citations/36413210), PubMed: [38355798](http://www.uniprot.org/citations/38355798)). In the activated complex, electrons are first transferred from NADPH to flavin adenine dinucleotide (FAD) and subsequently transferred via two heme molecules to molecular oxygen, producing superoxide through an outer-sphere reaction (Probable) (PubMed: [38355798](http://www.uniprot.org/citations/38355798)). Activation of the NADPH oxidase complex is initiated by the assembly of cytosolic subunits of the NADPH oxidase complex with the core NADPH oxidase complex to form a complex at the plasma

membrane or phagosomal membrane (PubMed:19028840, PubMed:38355798). This activation process is initiated by phosphorylation dependent binding of the cytosolic NCF1/p47-phox subunit to the C-terminus of CYBA/p22-phox (By similarity). NADPH oxidase complex assembly is impaired through interaction with NRROS (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein. Note=As unassembled monomer may localize to the endoplasmic reticulum

Tissue Location

Detected in neutrophils (at protein level).

gp91 phox Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

gp91 phox Antibody - Images**gp91 phox Antibody - Background**

Critical component of the membrane-bound oxidase of phagocytes that generates superoxide. It is the terminal component of a respiratory chain that transfers single electrons from cytoplasmic NADPH across the plasma membrane to molecular oxygen on the exterior. Also functions as a voltage-gated proton channel that mediates the H(+) currents of resting phagocytes. It participates in the regulation of cellular pH and is blocked by zinc.

gp91 phox Antibody - References

Royer-Pokora B.,et al.Nature 322:32-38(1986).
Jirapongsananuruk O.,et al.Clin. Immunol. 104:73-76(2002).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Dinauer M.C.,et al.Nature 327:717-720(1987).