

Desmin Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51162**Specification**

Desmin Antibody - Product Information

Application	WB, ICC, IHC-P, E
Primary Accession	P17661
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53 KDa

Desmin Antibody - Additional Information**Gene ID** 1674**Other Names**

Desmin, DES

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Desmin Antibody - Protein Information**Name** DES**Function**

Muscle-specific type III intermediate filament essential for proper muscular structure and function. Plays a crucial role in maintaining the structure of sarcomeres, inter-connecting the Z-disks and forming the myofibrils, linking them not only to the sarcolemmal cytoskeleton, but also to the nucleus and mitochondria, thus providing strength for the muscle fiber during activity (PubMed: [25358400](http://www.uniprot.org/citations/25358400)). In adult striated muscle they form a fibrous network connecting myofibrils to each other and to the plasma membrane from the periphery of the Z- line structures (PubMed: [24200904](http://www.uniprot.org/citations/24200904), PubMed: [25394388](http://www.uniprot.org/citations/25394388), PubMed: [26724190](http://www.uniprot.org/citations/26724190)). May act as a sarcomeric microtubule-anchoring protein: specifically associates with detyrosinated tubulin-alpha chains, leading to buckled microtubules and mechanical resistance to contraction. Required for nuclear membrane integrity, via anchoring at the cell tip and nuclear envelope, resulting in maintenance of microtubule-derived intracellular mechanical forces (By similarity). Contributes to the transcriptional regulation of the NKX2-5 gene in cardiac progenitor cells during a short period of cardiomyogenesis and in cardiac side population stem cells in the adult. Plays a role in maintaining an optimal conformation of nebulin (NEB) on heart muscle sarcomeres to bind and

recruit cardiac alpha-actin (By similarity).

Cellular Location

Cytoplasm, myofibril, sarcomere, Z line. Cytoplasm Cell membrane, sarcolemma. Nucleus {ECO:0000250|UniProtKB:P31001}. Cell tip {ECO:0000250|UniProtKB:P31001}. Nucleus envelope {ECO:0000250|UniProtKB:P31001}. Note=Localizes in the intercalated disks which occur at the Z line of cardiomyocytes (PubMed:24200904, PubMed:26724190). Localizes in the nucleus exclusively in differentiating cardiac progenitor cells and premature cardiomyocytes (By similarity). PKP2 is required for correct anchoring of DES at the cell tip and nuclear envelope (By similarity) {ECO:0000250|UniProtKB:P31001, ECO:0000269|PubMed:24200904, ECO:0000269|PubMed:26724190}

Desmin Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Desmin Antibody - Images

Desmin Antibody - Background

Desmin are class-III intermediate filaments found in muscle cells. In adult striated muscle they form a fibrous network connecting myofibrils to each other and to the plasma membrane from the periphery of the Z-line structures.

Desmin Antibody - References

Li Z.,et al.Gene 78:243-254(1989).
Li Z.,et al.J. Biol. Chem. 266:6562-6570(1991).
Vicart P.,et al.Hum. Genet. 98:422-429(1996).
Goldfarb L.G.,et al.Nat. Genet. 19:402-403(1998).
Li D.,et al.Circulation 100:461-464(1999).