

Histone H4 (AcK12) Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51256

Specification

Histone H4 (AcK12) Antibody - Product Information

Application	WB
Primary Accession	P62805
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	11 KDa
Antigen Region	1 - 60

Histone H4 (AcK12) Antibody - Additional Information

Gene ID 121504;554313;8294;8359;8360;8361;8362;8363;8364;8365;8366;8367;8368;8370

Other Names

Histone H4, HIST1H4A, H4/A, H4FA

Target/Specificity

KLH conjugated synthetic peptide derived from human Histone H4 (AcK12)

Dilution

WB~~ 1:1000

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Histone H4 (AcK12) Antibody - Protein Information

Name H4C1

Synonyms H4/A, H4FA, HIST1H4A

Function

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Cellular Location


Nucleus. Chromosome.

Histone H4 (AcK12) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Histone H4 (AcK12) Antibody - Images



All lanes : Anti-Histone H4 (AcK12) Antibody at 1:1000 dilution Lane 1: HeLa whole cell lysates
Lane 2: HepG2 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 11 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

Histone H4 (AcK12) Antibody - Background

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Histone H4 (AcK12) Antibody - References

- Sierra F., et al. *Nucleic Acids Res.* 11:7069-7086(1983).
Pauli U., et al. *Science* 236:1308-1311(1987).
Albig W., et al. *Genomics* 10:940-948(1991).
Drabent B., et al. *DNA Cell Biol.* 14:591-597(1995).
Albig W., et al. *Gene* 184:141-148(1997).