

5-HT1A Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51274

Specification

5-HT1A Antibody - Product Information

Application	WB
Primary Accession	P08908
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46 KDa
Antigen Region	291 - 350

5-HT1A Antibody - Additional Information

Gene ID 3350

Other Names

5-hydroxytryptamine receptor 1A, 5-HT-1A, 5-HT1A, G-21, Serotonin receptor 1A, HTR1A, ADRB2RL1, ADRBRL1

Target/Specificity

KLH conjugated synthetic peptide derived from human 5-HT1A

Dilution

WB~~ 1:1000

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

5-HT1A Antibody - Protein Information

Name HTR1A

Synonyms ADRB2RL1, ADRBRL1

Function

G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also functions as a receptor for various drugs and psychoactive substances. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase. Beta-arrestin family members inhibit signaling via G proteins and mediate activation of alternative signaling pathways. Signaling inhibits adenylate cyclase activity and activates a phosphatidylinositol-calcium second messenger system that regulates the release of Ca(2+) ions from intracellular stores. Plays a role in the

regulation of 5-hydroxytryptamine release and in the regulation of dopamine and 5-hydroxytryptamine metabolism. Plays a role in the regulation of dopamine and 5-hydroxytryptamine levels in the brain, and thereby affects neural activity, mood and behavior. Plays a role in the response to angiogenic stimuli.

Cellular Location

Cell membrane; Multi-pass membrane protein. Cell projection, dendrite
{ECO:0000250|UniProtKB:P19327}

Tissue Location


Detected in lymph nodes, thymus and spleen. Detected in activated T-cells, but not in resting T-cells

5-HT1A Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

5-HT1A Antibody - Images



Anti-5-HT1A Antibody at 1:1000 dilution + human brain lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 46 kDa Blocking/Dilution buffer: 5% NFD/MTBST.

5-HT1A Antibody - Background

G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also functions as a receptor for various drugs and psychoactive substances. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of

down-stream effectors, such as adenylate cyclase. Beta-arrestin family members inhibit signaling via G proteins and mediate activation of alternative signaling pathways. Signaling inhibits adenylate cyclase activity and activates a phosphatidylinositol-calcium second messenger system that regulates the release of Ca(2+) ions from intracellular stores. Plays a role in the regulation of 5-hydroxytryptamine release and in the regulation of dopamine and 5-hydroxytryptamine metabolism. Plays a role in the regulation of dopamine and 5-hydroxytryptamine levels in the brain, and thereby affects neural activity, mood and behavior. Plays a role in the response to anxiogenic stimuli.

5-HT1A Antibody - References

- Kobilka B.K.,et al.Nature 329:75-79(1987).
Saltzman A.G.,et al.Submitted (FEB-1991) to the EMBL/GenBank/DDBJ databases.
Levy F.O.,et al.Submitted (MAY-1992) to the EMBL/GenBank/DDBJ databases.
Kitano T.,et al.Mol. Biol. Evol. 21:936-944(2004).
Puhl H.L. III,et al.Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases.