

MLH1 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51343**Specification**

MLH1 Antibody - Product Information

Application	WB, ICC, E
Primary Accession	P40692
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	85 KDa

MLH1 Antibody - Additional Information**Gene ID** 4292**Other Names**

DNA mismatch repair protein Mlh1, MutL protein homolog 1, MLH1, COCA2

Dilution

WB~~1:1000

ICC~~N/A

E~~N/A

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

MLH1 Antibody - Protein Information**Name** MLH1**Synonyms** COCA2**Function**

Heterodimerizes with PMS2 to form MutL alpha, a component of the post-replicative DNA mismatch repair system (MMR). DNA repair is initiated by MutS alpha (MSH2-MSH6) or MutS beta (MSH2-MSH3) binding to a dsDNA mismatch, then MutL alpha is recruited to the heteroduplex. Assembly of the MutL-MutS-heteroduplex ternary complex in presence of RFC and PCNA is sufficient to activate endonuclease activity of PMS2. It introduces single-strand breaks near the mismatch and thus generates new entry points for the exonuclease EXO1 to degrade the strand containing the mismatch. DNA methylation would prevent cleavage and therefore assure that only the newly mutated DNA strand is going to be corrected. MutL alpha (MLH1-PMS2) interacts physically with the clamp loader subunits of DNA polymerase III, suggesting that it may play a role to recruit the DNA polymerase III to the site of the MMR. Also implicated in DNA damage signaling,

a process which induces cell cycle arrest and can lead to apoptosis in case of major DNA damages. Heterodimerizes with MLH3 to form MutL gamma which plays a role in meiosis.

Cellular Location

Nucleus. Chromosome. Note=Recruited to chromatin in a MCM9- dependent manner.

Tissue Location

Colon, lymphocytes, breast, lung, spleen, testis, prostate, thyroid, gall bladder and heart

MLH1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MLH1 Antibody - Images**MLH1 Antibody - Background**

Heterodimerizes with PMS2 to form MutL alpha, a component of the post-replicative DNA mismatch repair system (MMR). DNA repair is initiated by MutS alpha (MSH2-MSH6) or MutS beta (MSH2-MSH6) binding to a dsDNA mismatch, then MutL alpha is recruited to the heteroduplex. Assembly of the MutL-MutS- heteroduplex ternary complex in presence of RFC and PCNA is sufficient to activate endonuclease activity of PMS2. It introduces single-strand breaks near the mismatch and thus generates new entry points for the exonuclease EXO1 to degrade the strand containing the mismatch. DNA methylation would prevent cleavage and therefore assure that only the newly mutated DNA strand is going to be corrected. MutL alpha (MLH1-PMS2) interacts physically with the clamp loader subunits of DNA polymerase III, suggesting that it may play a role to recruit the DNA polymerase III to the site of the MMR. Also implicated in DNA damage signaling, a process which induces cell cycle arrest and can lead to apoptosis in case of major DNA damages. Heterodimerizes with MLH3 to form MutL gamma which plays a role in meiosis.

MLH1 Antibody - References

Bronner C.E.,et al.Nature 368:258-261(1994).
Papadopoulos N.,et al.Science 263:1625-1629(1994).
Kolodner R.D.,et al.Cancer Res. 55:242-248(1995).
Han H.-J.,et al.Hum. Mol. Genet. 4:237-242(1995).
Ota T.,et al.Nat. Genet. 36:40-45(2004).