

**PPHLN1 Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP51442**

## Specification

---

### PPHLN1 Antibody - Product Information

Application	WB, ICC, IHC-P, E
Primary Accession	<a href="#">Q8NEY8</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	53, 50, 43, 37 KDa

### PPHLN1 Antibody - Additional Information

Gene ID 51535

#### Other Names

Periphilin-1, Gastric cancer antigen Ga50, PPHLN1

#### Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

#### Storage

Store at -20 °C. Stable for 12 months from date of receipt

### PPHLN1 Antibody - Protein Information

Name PPHLN1 ([HGNC:19369](#))

#### Function

Component of the HUSH complex, a multiprotein complex that mediates epigenetic repression. The HUSH complex is recruited to genomic loci rich in H3K9me3 and is probably required to maintain transcriptional silencing by promoting recruitment of SETDB1, a histone methyltransferase that mediates further deposition of H3K9me3. In the HUSH complex, contributes to the maintenance of the complex at chromatin (PubMed: [26022416](http://www.uniprot.org/citations/26022416)). Acts as a transcriptional corepressor and regulates the cell cycle, probably via the HUSH complex (PubMed: [15474462](http://www.uniprot.org/citations/15474462), PubMed: [17963697](http://www.uniprot.org/citations/17963697)). The HUSH complex is also involved in the silencing of unintegrated retroviral DNA: some part of the retroviral DNA formed immediately after infection remains unintegrated in the host genome and is transcriptionally repressed (PubMed: [30487602](http://www.uniprot.org/citations/30487602)). May be involved in epithelial differentiation by contributing to epidermal integrity and barrier formation (PubMed: [12853457](http://www.uniprot.org/citations/12853457)).

#### Cellular Location

Nucleus. Cytoplasm. Chromosome. Note=In undifferentiated keratinocytes expressed in speckle-type nuclear granules and at the nuclear membrane, but in the differentiated keratinocytes colocalized with periplakin at the cell periphery and at cell-cell junctions (PubMed:12853457) Localizes to chromatin (PubMed:26022416).

**Tissue Location**

Ubiquitous..

**PPHLN1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**PPHLN1 Antibody - Images****PPHLN1 Antibody - Background**

Involved in epithelial differentiation and contributes to epidermal integrity and barrier formation.

**PPHLN1 Antibody - References**

Line A.,et al.Br. J. Cancer 86:1824-1830(2002).  
Kazerounian S.,et al.J. Biol. Chem. 278:36707-36717(2003).  
Zhang Q.-H.,et al.Genome Res. 10:1546-1560(2000).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Scherer S.E.,et al.Nature 440:346-351(2006).