

**SAP155 Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP51507**

**Specification**

---

**SAP155 Antibody - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">O75533</a>
Reactivity	<b>Human, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>155 KDa</b>
Antigen Region	<b>71 - 130</b>

**SAP155 Antibody - Additional Information**

**Gene ID** 23451

**Other Names**

Splicing factor 3B subunit 1, Pre-mRNA-splicing factor SF3b 155 kDa subunit, SF3b155, Spliceosome-associated protein 155, SAP 155, SF3B1, SAP155

**Target/Specificity**

KLH conjugated synthetic peptide derived from human SAP155

**Dilution**

WB~~ 1:1000

**Format**

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**SAP155 Antibody - Protein Information**

**Name** SF3B1 {ECO:0000303|PubMed:30567737, ECO:0000312|HGNC:HGNC:10768}

**Function**

Component of the 17S U2 SnRNP complex of the spliceosome, a large ribonucleoprotein complex that removes introns from transcribed pre-mRNAs (PubMed:<a href="http://www.uniprot.org/citations/12234937" target="\_blank">12234937</a>, PubMed:<a href="http://www.uniprot.org/citations/27720643" target="\_blank">27720643</a>, PubMed:<a href="http://www.uniprot.org/citations/32494006" target="\_blank">32494006</a>, PubMed:<a href="http://www.uniprot.org/citations/34822310" target="\_blank">34822310</a>). The 17S U2 SnRNP complex (1) directly participates in early spliceosome assembly and (2) mediates recognition of the intron branch site during pre-mRNA splicing by promoting the selection of the pre-mRNA branch-site adenosine, the nucleophile for the first step of splicing (PubMed:<a

<http://www.uniprot.org/citations/32494006> target="\_blank">32494006</a>, PubMed:<a href="http://www.uniprot.org/citations/34822310" target="\_blank">34822310</a>). Within the 17S U2 snRNP complex, SF3B1 is part of the SF3B subcomplex, which is required for 'A' complex assembly formed by the stable binding of U2 snRNP to the branchpoint sequence in pre-mRNA (PubMed:<a href="http://www.uniprot.org/citations/12234937" target="\_blank">12234937</a>). Sequence independent binding of SF3A and SF3B subcomplexes upstream of the branch site is essential, it may anchor U2 snRNP to the pre-mRNA (PubMed:<a href="http://www.uniprot.org/citations/12234937" target="\_blank">12234937</a>). May also be involved in the assembly of the 'E' complex (PubMed:<a href="http://www.uniprot.org/citations/10882114" target="\_blank">10882114</a>). Also acts as a component of the minor spliceosome, which is involved in the splicing of U12-type introns in pre-mRNAs (PubMed:<a href="http://www.uniprot.org/citations/15146077" target="\_blank">15146077</a>, PubMed:<a href="http://www.uniprot.org/citations/33509932" target="\_blank">33509932</a>). Together with other U2 snRNP complex components may also play a role in the selective processing of microRNAs (miRNAs) from the long primary miRNA transcript, pri-miR-17-92 (By similarity).

### Cellular Location


Nucleus. Nucleus speckle. Note=During mitosis, transiently dispersed from the nuclear speckles to the cytoplasm

### SAP155 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### SAP155 Antibody - Images



All lanes : Anti-SAP155 Antibody at 1:1000 dilution Lane 1: Hela whole cell lysates Lane 2: Ramos

whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 146 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

### **SAP155 Antibody - Background**

Subunit of the splicing factor SF3B required for 'A' complex assembly formed by the stable binding of U2 snRNP to the branchpoint sequence (BPS) in pre-mRNA. Sequence independent binding of SF3A/SF3B complex upstream of the branch site is essential, it may anchor U2 snRNP to the pre-mRNA. May also be involved in the assembly of the 'E' complex. Belongs also to the minor U12-dependent spliceosome, which is involved in the splicing of rare class of nuclear pre-mRNA intron.

### **SAP155 Antibody - References**

Wang C., et al. Genes Dev. 12:1409-1414(1998).  
Woessner J., et al. Submitted (SEP-1998) to the EMBL/GenBank/DDBJ databases.  
Hillier L.W., et al. Nature 434:724-731(2005).  
Yu W., et al. Submitted (JUN-1998) to the EMBL/GenBank/DDBJ databases.  
Das R., et al. Mol. Cell 5:779-787(2000).