

CD184 (pS339) Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51641

Specification

CD184 (pS339) Antibody - Product Information

Application	WB
Primary Accession	P61073
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47 KDa

CD184 (pS339) Antibody - Additional Information

Gene ID 7852

Other Names

C-X-C chemokine receptor type 4, CXC-R4, CXCR-4, FB22, Fusin, HM89, LCR1, Leukocyte-derived seven transmembrane domain receptor, LESTR, Lipopolysaccharide-associated protein 3, LAP-3, LPS-associated protein 3, NPYRL, Stromal cell-derived factor 1 receptor, SDF-1 receptor, CD184, CXCR4

Dilution

WB~~1:500

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

CD184 (pS339) Antibody - Protein Information

Name CXCR4

Function

Receptor for the C-X-C chemokine CXCL12/SDF-1 that transduces a signal by increasing intracellular calcium ion levels and enhancing MAPK1/MAPK3 activation (PubMed: [10452968](http://www.uniprot.org/citations/10452968), PubMed: [18799424](http://www.uniprot.org/citations/18799424), PubMed: [24912431](http://www.uniprot.org/citations/24912431), PubMed: [28978524](http://www.uniprot.org/citations/28978524)). Involved in the AKT signaling cascade (PubMed: [24912431](http://www.uniprot.org/citations/24912431)). Plays a role in regulation of cell migration, e.g. during wound healing (PubMed: [28978524](http://www.uniprot.org/citations/28978524)). Acts as a receptor for extracellular ubiquitin; leading to enhanced intracellular calcium ions and reduced cellular cAMP levels (PubMed: [28978524](http://www.uniprot.org/citations/28978524)).

<http://www.uniprot.org/citations/20228059> target="_blank">20228059). Binds bacterial lipopolysaccharide (LPS) et mediates LPS-induced inflammatory response, including TNF secretion by monocytes (PubMed:11276205). Involved in hematopoiesis and in cardiac ventricular septum formation. Also plays an essential role in vascularization of the gastrointestinal tract, probably by regulating vascular branching and/or remodeling processes in endothelial cells. Involved in cerebellar development. In the CNS, could mediate hippocampal-neuron survival (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein. Cell junction. Early endosome. Late endosome. Lysosome. Note=In unstimulated cells, diffuse pattern on plasma membrane. On agonist stimulation, colocalizes with ITCH at the plasma membrane where it becomes ubiquitinated. In the presence of antigen, distributes to the immunological synapse forming at the T- cell-APC contact area, where it localizes at the peripheral and distal supramolecular activation cluster (SMAC)

Tissue Location


Expressed in numerous tissues, such as peripheral blood leukocytes, spleen, thymus, spinal cord, heart, placenta, lung, liver, skeletal muscle, kidney, pancreas, cerebellum, cerebral cortex and medulla (in microglia as well as in astrocytes), brain microvascular, coronary artery and umbilical cord endothelial cells Isoform 1 is predominant in all tissues tested

CD184 (pS339) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD184 (pS339) Antibody - Images



CD184 (pS339) Antibody - Background

Receptor for the C-X-C chemokine CXCL12/SDF-1 that transduces a signal by increasing intracellular calcium ion levels and enhancing MAPK1/MAPK3 activation. Acts as a receptor for extracellular ubiquitin; leading to enhanced intracellular calcium ions and reduced cellular cAMP levels. Involved in hematopoiesis and in cardiac ventricular septum formation. Also plays an essential role in vascularization of the gastrointestinal tract, probably by regulating vascular branching and/or remodeling processes in endothelial cells. Involved in cerebellar development. In the CNS, could mediate hippocampal-neuron survival. Acts as a coreceptor (CD4 being the primary receptor) for HIV-1 X4 isolates and as a primary receptor for some HIV-2 isolates. Promotes Env-mediated fusion of the virus.

CD184 (pS339) Antibody - References

Herzog H.,et al.DNA Cell Biol. 12:465-471(1993).
Jazin E.E.,et al.Regul. Pept. 47:247-258(1993).
Federspiel B.,et al.Genomics 16:707-712(1993).
Loetscher M.,et al.J. Biol. Chem. 269:232-237(1994).
Nomura H.,et al.Int. Immunol. 5:1239-1249(1993).