

NDC80 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51659**Specification**

NDC80 Antibody - Product Information

Application	WB, E
Primary Accession	O14777
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	74 KDa

NDC80 Antibody - Additional Information**Gene ID** 10403**Other Names**

Kinetochores protein NDC80 homolog, Highly expressed in cancer protein, Kinetochores protein Hec1, HsHec1, Kinetochores-associated protein 2, Retinoblastoma-associated protein HEC, NDC80, HEC, HEC1, KNTC2

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

NDC80 Antibody - Protein Information**Name** NDC80**Synonyms** HEC, HEC1, KNTC2**Function**

Acts as a component of the essential kinetochores-associated NDC80 complex, which is required for chromosome segregation and spindle checkpoint activity (PubMed: [12351790](http://www.uniprot.org/citations/12351790), PubMed: [14654001](http://www.uniprot.org/citations/14654001), PubMed: [14699129](http://www.uniprot.org/citations/14699129), PubMed: [15062103](http://www.uniprot.org/citations/15062103), PubMed: [15235793](http://www.uniprot.org/citations/15235793), PubMed: [15239953](http://www.uniprot.org/citations/15239953), PubMed: [15548592](http://www.uniprot.org/citations/15548592), PubMed: [16732327](http://www.uniprot.org/citations/16732327), PubMed: [30409912](http://www.uniprot.org/citations/30409912), PubMed: [9315664](http://www.uniprot.org/citations/9315664)). Required for kinetochores integrity and the organization of stable microtubule binding sites in the outer plate of

the kinetochore (PubMed:15548592, PubMed:30409912). The NDC80 complex synergistically enhances the affinity of the SKA1 complex for microtubules and may allow the NDC80 complex to track depolymerizing microtubules (PubMed:23085020). Plays a role in chromosome congression and is essential for the end-on attachment of the kinetochores to spindle microtubules (PubMed:23891108, PubMed:25743205).

Cellular Location

Nucleus. Chromosome, centromere, kinetochore. Note=Localizes to kinetochores from late prophase to anaphase (PubMed:14699129) Localizes specifically to the outer plate of the kinetochore (PubMed:14699129).

NDC80 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NDC80 Antibody - Images

NDC80 Antibody - Background

Acts as a component of the essential kinetochore-associated NDC80 complex, which is required for chromosome segregation and spindle checkpoint activity. Required for kinetochore integrity and the organization of stable microtubule binding sites in the outer plate of the kinetochore.

NDC80 Antibody - References

Chen Y.,et al.Mol. Cell. Biol. 17:6049-6056(1997).
Chen Y.,et al.J. Biol. Chem. 272:24081-24087(1997).
Zheng L.,et al.Mol. Cell. Biol. 19:5417-5428(1999).
Zheng L.,et al.Mol. Cell. Biol. 20:3529-3537(2000).
Chen Y.,et al.J. Biol. Chem. 277:49408-49416(2002).