

Fibulin 1 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51770**Specification**

Fibulin 1 Antibody - Product Information

Application	WB, E
Primary Accession	P23142
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	77 KDa

Fibulin 1 Antibody - Additional Information**Gene ID** 2192**Other Names**

Fibulin-1, FIBL-1, FBLN1

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Fibulin 1 Antibody - Protein Information**Name** FBLN1**Function**

Incorporated into fibronectin-containing matrix fibers. May play a role in cell adhesion and migration along protein fibers within the extracellular matrix (ECM). Could be important for certain developmental processes and contribute to the supramolecular organization of ECM architecture, in particular to those of basement membranes. Has been implicated in a role in cellular transformation and tumor invasion, it appears to be a tumor suppressor. May play a role in haemostasis and thrombosis owing to its ability to bind fibrinogen and incorporate into clots. Could play a significant role in modulating the neurotrophic activities of APP, particularly soluble APP.

Cellular Location

Secreted, extracellular space, extracellular matrix

Tissue Location

Isoform A and isoform B are only expressed in placenta. Isoform C and isoform D are expressed in a variety of tissues and cultured cells.

Fibulin 1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Fibulin 1 Antibody - Images

Fibulin 1 Antibody - Background

Incorporated into fibronectin-containing matrix fibers. May play a role in cell adhesion and migration along protein fibers within the extracellular matrix (ECM). Could be important for certain developmental processes and contribute to the supramolecular organization of ECM architecture, in particular to those of basement membranes. Has been implicated in a role in cellular transformation and tumor invasion, it appears to be a tumor suppressor. May play a role in haemostasis and thrombosis owing to its ability to bind fibrinogen and incorporate into clots. Could play a significant role in modulating the neurotrophic activities of APP, particularly soluble APP.

Fibulin 1 Antibody - References

Argraves W.S., et al. J. Cell Biol. 111:3155-3164(1990).
Tran H., et al. Matrix Biol. 15:479-493(1997).
Krichevsky A.M., et al. J. Biol. Chem. 274:14295-14305(1999).
Wan D., et al. Proc. Natl. Acad. Sci. U.S.A. 101:15724-15729(2004).
Dunham I., et al. Nature 402:489-495(1999).