

**Shc (Phospho-Tyr349) Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP52353**

## Specification

---

### Shc (Phospho-Tyr349) Antibody - Product Information

Application	WB, IHC
Primary Accession	<a href="#">P29353</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	62822

### Shc (Phospho-Tyr349) Antibody - Additional Information

**Gene ID** 6464

#### Other Names

SHC-transforming protein 1, SHC-transforming protein 3, SHC-transforming protein A, Src homology 2 domain-containing-transforming protein C1, SH2 domain protein C1, SHC1, SHC, SHCA

#### Dilution

WB~~1:1000  
IHC~~1:50~100

#### Format

Rabbit IgG in phosphate buffered saline (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

#### Storage Conditions

-20°C

### Shc (Phospho-Tyr349) Antibody - Protein Information

**Name** SHC1

**Synonyms** SHC, SHCA

#### Function

Signaling adapter that couples activated growth factor receptors to signaling pathways. Participates in a signaling cascade initiated by activated KIT and KITLG/SCF. Isoform p46Shc and isoform p52Shc, once phosphorylated, couple activated receptor tyrosine kinases to Ras via the recruitment of the GRB2/SOS complex and are implicated in the cytoplasmic propagation of mitogenic signals. Isoform p46Shc and isoform p52Shc may thus function as initiators of the Ras signaling cascade in various non-neuronal systems. Isoform p66Shc does not mediate Ras activation, but is involved in signal transduction pathways that regulate the cellular response to oxidative stress and life span. Isoform p66Shc acts as a downstream target of the tumor suppressor p53 and is indispensable for the ability of stress-activated p53 to induce elevation of

intracellular oxidants, cytochrome c release and apoptosis. The expression of isoform p66Shc has been correlated with life span (By similarity). Participates in signaling downstream of the angiopoietin receptor TEK/TIE2, and plays a role in the regulation of endothelial cell migration and sprouting angiogenesis.

#### Cellular Location

Cytoplasm. Cell junction, focal adhesion [Isoform p66Shc]: Mitochondrion. Note=In case of oxidative conditions, phosphorylation at 'Ser-36' of isoform p66Shc, leads to mitochondrial accumulation.

#### Tissue Location


Widely expressed. Expressed in neural stem cells but absent in mature neurons

### Shc (Phospho-Tyr349) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Shc (Phospho-Tyr349) Antibody - Images



Western blot analysis of extracts from 293 cells treated with EGF (200ng/ml, 30mins), using Shc (phospho-Tyr349) antibody.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Shc (phospho-Tyr349) antibody

### **Shc (Phospho-Tyr349) Antibody - Background**

Signaling adapter that couples activated growth factor receptors to signaling pathways. Participates in a signaling cascade initiated by activated KIT and KITLG/SCF. Isoform p46Shc and isoform p52Shc, once phosphorylated, couple activated receptor tyrosine kinases to Ras via the recruitment of the GRB2/SOS complex and are implicated in the cytoplasmic propagation of mitogenic signals. Isoform p46Shc and isoform p52Shc may thus function as initiators of the Ras signaling cascade in various non-neuronal systems. Isoform p66Shc does not mediate Ras activation, but is involved in signal transduction pathways that regulate the cellular response to oxidative stress and life span. Isoform p66Shc acts as a downstream target of the tumor suppressor p53 and is indispensable for the ability of stress-activated p53 to induce elevation of intracellular oxidants, cytochrome c release and apoptosis. The expression of isoform p66Shc has been correlated with life span (By similarity). Participates in signaling downstream of the angiotensin receptor TEK/TIE2, and plays a role in the regulation of endothelial cell migration and sprouting angiogenesis.

### **Shc (Phospho-Tyr349) Antibody - References**

Pellicci G., et al. Cell 70:93-104(1992).  
Migliaccio E., et al. EMBO J. 16:706-716(1997).  
Harun R.B., et al. Genomics 42:349-352(1997).  
Ota T., et al. Nat. Genet. 36:40-45(2004).  
Goshima N., et al. Nat. Methods 5:1011-1017(2008).