

**IRF-3 (Phospho-Ser385) Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP52385****Specification**

---

**IRF-3 (Phospho-Ser385) Antibody - Product Information**

Application	IF, WB, IHC
Primary Accession	<a href="#">Q14653</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

**IRF-3 (Phospho-Ser385) Antibody - Additional Information****Gene ID** 3661**Other Names**

Interferon regulatory factor 3, IRF-3, IRF3

**Dilution**

IF~~1:100

WB~~1:1000

IHC~~1:50~100

**Format**Rabbit IgG in phosphate buffered saline (without Mg<sup>2+</sup> and Ca<sup>2+</sup>), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.**Storage Conditions**

-20°C

**IRF-3 (Phospho-Ser385) Antibody - Protein Information****Name** IRF3 {ECO:0000303|PubMed:9803267, ECO:0000312|HGNC:HGNC:6118}**Function**

Key transcriptional regulator of type I interferon (IFN)- dependent immune responses which plays a critical role in the innate immune response against DNA and RNA viruses (PubMed:<a href="http://www.uniprot.org/citations/22394562" target="\_blank">22394562</a>, PubMed:<a href="http://www.uniprot.org/citations/24049179" target="\_blank">24049179</a>, PubMed:<a href="http://www.uniprot.org/citations/25636800" target="\_blank">25636800</a>, PubMed:<a href="http://www.uniprot.org/citations/27302953" target="\_blank">27302953</a>, PubMed:<a href="http://www.uniprot.org/citations/31340999" target="\_blank">31340999</a>, PubMed:<a href="http://www.uniprot.org/citations/36603579" target="\_blank">36603579</a>, PubMed:<a href="http://www.uniprot.org/citations/8524823" target="\_blank">8524823</a>). Regulates the transcription of type I IFN genes (IFN-alpha and IFN-beta) and IFN-stimulated genes (ISG) by binding to an interferon-stimulated response element (ISRE) in their promoters (PubMed:<a href="http://www.uniprot.org/citations/11846977" target="\_blank">11846977</a>, PubMed:<a href="http://www.uniprot.org/citations/11846977" target="\_blank">11846977</a>).

<http://www.uniprot.org/citations/16846591> target="\_blank">16846591</a>, PubMed:<a href="http://www.uniprot.org/citations/16979567" target="\_blank">16979567</a>, PubMed:<a href="http://www.uniprot.org/citations/20049431" target="\_blank">20049431</a>, PubMed:<a href="http://www.uniprot.org/citations/32972995" target="\_blank">32972995</a>, PubMed:<a href="http://www.uniprot.org/citations/36603579" target="\_blank">36603579</a>, PubMed:<a href="http://www.uniprot.org/citations/8524823" target="\_blank">8524823</a>). Acts as a more potent activator of the IFN-beta (IFNB) gene than the IFN-alpha (IFNA) gene and plays a critical role in both the early and late phases of the IFNA/B gene induction (PubMed:<a href="http://www.uniprot.org/citations/16846591" target="\_blank">16846591</a>, PubMed:<a href="http://www.uniprot.org/citations/16979567" target="\_blank">16979567</a>, PubMed:<a href="http://www.uniprot.org/citations/20049431" target="\_blank">20049431</a>, PubMed:<a href="http://www.uniprot.org/citations/36603579" target="\_blank">36603579</a>). Found in an inactive form in the cytoplasm of uninfected cells and following viral infection, double-stranded RNA (dsRNA), or toll-like receptor (TLR) signaling, is phosphorylated by IKBKE and TBK1 kinases (PubMed:<a href="http://www.uniprot.org/citations/22394562" target="\_blank">22394562</a>, PubMed:<a href="http://www.uniprot.org/citations/25636800" target="\_blank">25636800</a>, PubMed:<a href="http://www.uniprot.org/citations/27302953" target="\_blank">27302953</a>, PubMed:<a href="http://www.uniprot.org/citations/36603579" target="\_blank">36603579</a>). This induces a conformational change, leading to its dimerization and nuclear localization and association with CREB binding protein (CREBBP) to form dsRNA-activated factor 1 (DRAF1), a complex which activates the transcription of the type I IFN and ISG genes (PubMed:<a href="http://www.uniprot.org/citations/16154084" target="\_blank">16154084</a>, PubMed:<a href="http://www.uniprot.org/citations/27302953" target="\_blank">27302953</a>, PubMed:<a href="http://www.uniprot.org/citations/33440148" target="\_blank">33440148</a>, PubMed:<a href="http://www.uniprot.org/citations/36603579" target="\_blank">36603579</a>). Can activate distinct gene expression programs in macrophages and can induce significant apoptosis in primary macrophages (PubMed:<a href="http://www.uniprot.org/citations/16846591" target="\_blank">16846591</a>). In response to Sendai virus infection, is recruited by TOMM70:HSP90AA1 to mitochondrion and forms an apoptosis complex TOMM70:HSP90AA1:IRF3:BAX inducing apoptosis (PubMed:<a href="http://www.uniprot.org/citations/25609812" target="\_blank">25609812</a>). Key transcription factor regulating the IFN response during SARS-CoV-2 infection (PubMed:<a href="http://www.uniprot.org/citations/33440148" target="\_blank">33440148</a>).

### Cellular Location

Cytoplasm. Nucleus Mitochondrion. Note=Shuttles between cytoplasmic and nuclear compartments, with export being the prevailing effect (PubMed:10805757, PubMed:35922005). When activated, IRF3 interaction with CREBBP prevents its export to the cytoplasm (PubMed:10805757). Recruited to mitochondria via TOMM70:HSP90AA1 upon Sendai virus infection (PubMed:25609812).

### Tissue Location

Expressed constitutively in a variety of tissues.


## IRF-3 (Phospho-Ser385) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)


### IRF-3 (Phospho-Ser385) Antibody - Images



Immunofluorescence analysis of HeLa cells, using IRF-3 (Phospho-Ser385) antibody.



Western blot analysis of extracts from HT-29 cells, treated with insulin (0.01U/ml, 15mins), using IRF-3 (Phospho-Ser385) antibody.



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue using IRF-3 (Phospho-Ser385) antibody.

### IRF-3 (Phospho-Ser385) Antibody - Background

Key transcriptional regulator of type I interferon (IFN)-dependent immune responses which plays a critical role in the innate immune response against DNA and RNA viruses. Regulates the transcription of type I IFN genes (IFN-alpha and IFN-beta) and IFN-stimulated genes (ISG) by binding to an interferon-stimulated response element (ISRE) in their promoters. Acts as a more potent activator of the IFN-beta (IFNB) gene than the IFN-alpha (IFNA) gene and plays a critical role in both the early and late phases of the IFNA/B gene induction. Found in an inactive form in the cytoplasm of uninfected cells and following viral infection, double-stranded RNA (dsRNA), or toll-like receptor

(TLR) signaling, is phosphorylated by IKKε and TBK1 kinases. This induces a conformational change, leading to its dimerization and nuclear localization and association with CREB binding protein (CREBBP) to form dsRNA-activated factor 1 (DAF1), a complex which activates the transcription of the type I IFN and ISG genes. Can activate distinct gene expression programs in macrophages and can induce significant apoptosis in primary macrophages.

### **IRF-3 (Phospho-Ser385) Antibody - References**

Au W.W.-C., et al. Proc. Natl. Acad. Sci. U.S.A. 92:11657-11661(1995).  
Tabata Y., et al. Submitted (FEB-2003) to the EMBL/GenBank/DDBJ databases.  
Ota T., et al. Nat. Genet. 36:40-45(2004).  
Grimwood J., et al. Nature 428:529-535(2004).  
Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.