

**KLF4 Antibody**  
**Purified Mouse Monoclonal Antibody (Mab)**  
**Catalog # AP52656**

**Specification**

---

**KLF4 Antibody - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">O43474</a>
Reactivity	<b>Transfected</b>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>
Isotype	<b>IgG2b</b>
Calculated MW	<b>55/65 KDa</b>

**KLF4 Antibody - Additional Information**

**Gene ID** 9314

**Other Names**

Endothelial Kruppel like zinc finger protein; Epithelial zinc finger protein  
EZf; EZF; GKLF; Gut-enriched krueppel-like factor; KLF; KLF4; KLF4\_HUMAN; Krueppel-like factor  
4; Kruppel like factor 4 (Epithelial zinc finger protein EZF) (Gut enriched Krueppel like  
factor); Kruppel like factor 4 (gut).

**Dilution**

WB ~ ~ 1:1000

**Format**

ascites

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**KLF4 Antibody - Protein Information**

**Name** KLF4 ([HGNC:6348](#))

**Synonyms** EZF, GKLF

**Function**

Transcription factor; can act both as activator and as repressor. Binds the 5'-CACCC-3' core sequence. Binds to the promoter region of its own gene and can activate its own transcription. Regulates the expression of key transcription factors during embryonic development. Plays an important role in maintaining embryonic stem cells, and in preventing their differentiation. Required for establishing the barrier function of the skin and for postnatal maturation and maintenance of the ocular surface. Involved in the differentiation of epithelial cells and may also function in skeletal and kidney development. Contributes to the down-regulation of p53/TP53 transcription.

### Cellular Location


Nucleus {ECO:0000250|UniProtKB:Q60793}. Cytoplasm {ECO:0000250|UniProtKB:Q60793}

### KLF4 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### KLF4 Antibody - Images



Western blot analysis of extracts from CHO-K1 cells, transfected with pcDNA3.1-Hygro(+)-mKLF4-Flag construct (A) or transfected with pDNA3.1-Hygro(+)-Flag vector (B), using KLF4 mouse mAb (1:1000 diluted). Predicted band size: 55/65 kDa. Observed band size: 55/65 kDa.

### KLF4 Antibody - Background

Transcription factor; can act both as activator and as repressor. Binds the 5'-CACCC-3' core sequence. Binds to the promoter region of its own gene and can activate its own transcription. Regulates the expression of key transcription factors during embryonic development. Plays an important role in maintaining embryonic stem cells, and in preventing their differentiation. Required for establishing the barrier function of the skin and for postnatal maturation and maintenance of the ocular surface. Involved in the differentiation of epithelial cells and may also function in skeletal and kidney development. Contributes to the down-regulation of p53/TP53 transcription.

### KLF4 Antibody - References

Yet S.-F., et al. J. Biol. Chem. 273:1026-1031(1998).  
Foster K.W., et al. Cell Growth Differ. 10:423-434(1999).

Camacho-Vanegas O., et al. *FASEB J.* 27:432-436(2013).

Garrett-Sinha L.A., et al. Submitted (SEP-1996) to the EMBL/GenBank/DDBJ databases.

Ota T., et al. *Nat. Genet.* 36:40-45(2004).