

COX IV Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AP52659**Specification**

COX IV Antibody - Product Information

Application	IP, WB, FC, ICC
Primary Accession	P13073
Reactivity	Human, Mouse, Neisseria Gonorrhoeae
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	17 KDa

COX IV Antibody - Additional Information**Gene ID** 1327**Other Names**

AL024441;COX 4;COX IV 1;COX IV;COX IV-1;Cox4;COX41_HUMAN;Cox4a;COX4B;COX4I1;COX4I2;COX4L2;COXIV;Cytochrome c oxidase polypeptide IV;Cytochrome c oxidase subunit 4 isoform 1 mitochondrial;Cytochrome c oxidase subunit 4 isoform 1, mitochondrial;Cytochrome C Oxidase subunit IV;Cytochrome c oxidase subunit IV isoform 1;Cytochrome c oxidase subunit IV isoform 2 (lung);Cytochrome c oxydase subunit;dj857M17.2;MGC105470;MGC72016.

Dilution

IP~~1:500
WB~~1:1000
FC~~1:100
ICC~~1:150

Format

Purified mouse monoclonal antibody in buffer containing 0.1M Tris-Glycine(pH 7.4,150 mM NaCl)with 0.09% (W/V) sodium azide,0.1mg/mlBSA and 50% glycerol.

Storage

Store at -20 °C.Stable for 12 months from date of receipt

COX IV Antibody - Protein Information**Name** COX4I1 ([HGNC:2265](#))**Function**

Component of the cytochrome c oxidase, the last enzyme in the mitochondrial electron transport chain which drives oxidative phosphorylation. The respiratory chain contains 3 multisubunit complexes succinate dehydrogenase (complex II, CII), ubiquinol- cytochrome c oxidoreductase (cytochrome b-c1 complex, complex III, CIII) and cytochrome c oxidase (complex IV, CIV), that cooperate to transfer electrons derived from NADH and succinate to molecular oxygen, creating an

electrochemical gradient over the inner membrane that drives transmembrane transport and the ATP synthase. Cytochrome c oxidase is the component of the respiratory chain that catalyzes the reduction of oxygen to water. Electrons originating from reduced cytochrome c in the intermembrane space (IMS) are transferred via the dinuclear copper A center (CU(A)) of subunit 2 and heme A of subunit 1 to the active site in subunit 1, a binuclear center (BNC) formed by heme A3 and copper B (CU(B)). The BNC reduces molecular oxygen to 2 water molecules using 4 electrons from cytochrome c in the IMS and 4 protons from the mitochondrial matrix.

Cellular Location

Mitochondrion inner membrane; Single-pass membrane protein

Tissue Location


Ubiquitous.

COX IV Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


COX IV Antibody - Images




Immunoprecipitation analysis of HeLa cell lysates using COX IV mouse mAb.



Western blot detection of COX IV in Mouse skeletal muscle and Jurkat lysates using COX IV mouse mAb (1:1000 diluted). Predicted band size: 17KDa. Observed band size: 17KDa.



Flow Cytometry analysis of K562 cells stained with COX4 (red, 1/100 dilution), followed by FITC-conjugated goat anti-mouse IgG. Blue line histogram represents the isotype control, normal mouse IgG.



Immunocytochemistry of HeLa cells using anti-COX IV mouse mAb diluted 1:150.

COX IV Antibody - Background

This protein is one of the nuclear-coded polypeptide chains of cytochrome c oxidase, the terminal

oxidase in mitochondrial electron transport.

COX IV Antibody - References

Zeviani M., et al. Gene 55:205-217(1987).

Lomax M.I., et al. Gene 86:209-216(1990).

Park S.J., et al. Submitted (OCT-1990) to the EMBL/GenBank/DDBJ databases.

Yu W., et al. Submitted (MAR-1997) to the EMBL/GenBank/DDBJ databases.

Bachman N.J., et al. Submitted (MAY-1997) to the EMBL/GenBank/DDBJ databases.

COX IV Antibody - Citations

- [Mechanistic characterization of nitrite-mediated neuroprotection after experimental cardiac arrest.](#)