

Hsp70 (C-terminus) Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AP52677**Specification**

Hsp70 (C-terminus) Antibody - Product Information

Application	IP, WB
Primary Accession	P08107
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2b
Calculated MW	70 KDa

Hsp70 (C-terminus) Antibody - Additional Information**Other Names**

DAQB 147D11.1 001;FLJ54303;FLJ54370;FLJ54392;FLJ54408;FLJ75127;Heat shock 70 kDa protein 1;Heat shock 70 kDa protein 1/2;Heat shock 70 kDa protein 1A/1B;heat shock 70kDa protein 1A;Heat shock 70kDa protein 1B;Heat shock induced protein;heat shock protein 70;HSP70 1;HSP70 2;HSP70-1/HSP70-2;HSP70-1A;HSP70.1;HSP70.1/HSP70.2;HSP70I;HSP71_HUMAN;HSP72;HSPA1;HSPA1A;HSPA1B;XXbac BCX40G17.3 001.

Dilution

IP~~1:500
WB~~1:1000

Format

Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine(pH 7.4,150 mM NaCl)with 0.09% (W/V) sodium azide,0.1mg/mlBSA and 50% glycerol.

Storage

Store at -20 °C.Stable for 12 months from date of receipt


Hsp70 (C-terminus) Antibody - Protein Information**Hsp70 (C-terminus) Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

Hsp70 (C-terminus) Antibody - Images



Immunoprecipitation analysis of HeLa cell lysates using Hsp70 (C-terminus) mouse mAb.



Western blot detection of Hsp70(C-terminus) in HeLa and HsHeLa cell lysates using Hsp70(C-terminus) mouse mAb (1:1000 diluted). Predicted band size: 70KDa. Observed band size: 70KDa.

Hsp70 (C-terminus) Antibody - Background

In cooperation with other chaperones, Hsp70s stabilize preexistent proteins against aggregation and mediate the folding of newly translated polypeptides in the cytosol as well as within organelles. These chaperones participate in all these processes through their ability to recognize nonnative conformations of other proteins. They bind extended peptide segments with a net hydrophobic character exposed by polypeptides during translation and membrane translocation, or following stress-induced damage. In case of rotavirus A infection, serves as a post-attachment receptor for the virus to facilitate entry into the cell.

Hsp70 (C-terminus) Antibody - References

Milner C.M.,et al.Immunogenetics 32:242-251(1990).
Hunt C.,et al.Proc. Natl. Acad. Sci. U.S.A. 82:6455-6459(1985).
Xie T.,et al.Genome Res. 13:2621-2636(2003).
Shiina S.,et al.Submitted (SEP-1999) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).