

BiP/GRP78 (C-terminus) Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AP52683**Specification****BiP/GRP78 (C-terminus) Antibody - Product Information**

Application	WB, IHC, ICC
Primary Accession	P11021
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	78 KDa

BiP/GRP78 (C-terminus) Antibody - Additional Information**Gene ID** 3309**Other Names**

78 kDa glucose regulated protein;78 kDa glucose-regulated protein;AL022860;AU019543;BIP;D2Wsu141e;D2Wsu17e;Endoplasmic reticulum luminal Ca(2⁺)-binding protein grp78; Endoplasmic reticulum luminal Ca²⁺ binding protein grp78;FLJ26106;Glucose Regulated Protein 78kDa;GRP 78;GRP-78;GRP78;GRP78_HUMAN;Heat shock 70 kDa protein 5;Heat Shock 70kDa Protein 5;Hsce70;HSPA 5;HSPA5;Immunoglobulin Heavy Chain Binding Protein; Immunoglobulin heavy chain-binding protein;mBiP;MIF2;Sez7.

Dilution

WB~~1:1000

IHC~~1:100

ICC~~1:50

Format

Purified mouse monoclonal antibody in PBS(pH 7.4) containing with 0.09% (W/V) sodium azide and 50% glycerol.

Storage

Store at -20 °C.Stable for 12 months from date of receipt

BiP/GRP78 (C-terminus) Antibody - Protein Information**Name** HSPA5 ([HGNC:5238](#))**Function**

Endoplasmic reticulum chaperone that plays a key role in protein folding and quality control in the endoplasmic reticulum lumen (PubMed:2294010, PubMed:23769672, PubMed:23990668, PubMed:28332555

target="_blank">28332555). Involved in the correct folding of proteins and degradation of misfolded proteins via its interaction with DNAJC10/ERdj5, probably to facilitate the release of DNAJC10/ERdj5 from its substrate (By similarity). Acts as a key repressor of the ERN1/IRE1-mediated unfolded protein response (UPR) (PubMed:1550958, PubMed:19538957). In the unstressed endoplasmic reticulum, recruited by DNAJB9/ERdj4 to the luminal region of ERN1/IRE1, leading to disrupt the dimerization of ERN1/IRE1, thereby inactivating ERN1/IRE1 (By similarity). Accumulation of misfolded protein in the endoplasmic reticulum causes release of HSPA5/BiP from ERN1/IRE1, allowing homodimerization and subsequent activation of ERN1/IRE1 (By similarity). Plays an auxiliary role in post-translational transport of small presecretory proteins across endoplasmic reticulum (ER). May function as an allosteric modulator for SEC61 channel-forming translocon complex, likely cooperating with SEC62 to enable the productive insertion of these precursors into SEC61 channel. Appears to specifically regulate translocation of precursors having inhibitory residues in their mature region that weaken channel gating. May also play a role in apoptosis and cell proliferation (PubMed:26045166).

Cellular Location


Endoplasmic reticulum lumen. Melanosome. Cytoplasm {ECO:0000250|UniProtKB:P20029}. Cell surface Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:12643545). Localizes to the cell surface of epithelial cells in response to high levels of free iron (PubMed:20484814, PubMed:24355926, PubMed:27159390)


BiP/GRP78 (C-terminus) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BiP/GRP78 (C-terminus) Antibody - Images






Western blot detection of BiP/GRP78 (C-terminus) in HeLa, C6, Lncap and MDA-MB-468 cell lysates using BiP/GRP78 (C-terminus) mouse mAb (1:1000 diluted). Predicted band size: 72KDa. Observed band size: 78KDa.



Immunohistochemical analysis of paraffin-embedded Colorectal cancer using BiP/GRP78 (C-terminus) Mouse mAb (1/100 dilution). Antigen retrieval was performed by pressure cooking in citrate buffer (pH 6.0).



Immunocytochemistry staining of HeLa cells fixed with 4% Paraformaldehyde and using anti-BiP/GRP78 (C-terminus) mouse mAb (dilution 1:50).

BiP/GRP78 (C-terminus) Antibody - Background

Probably plays a role in facilitating the assembly of multimeric protein complexes inside the endoplasmic reticulum. Involved in the correct folding of proteins and degradation of misfolded proteins via its interaction with DNAJC10, probably to facilitate the release of DNAJC10 from its substrate.

BiP/GRP78 (C-terminus) Antibody - References

Ting J.,et al.DNA 7:275-286(1988).
Chao C.C.K.,et al.Submitted (DEC-1995) to the EMBL/GenBank/DDBJ databases.
Hansen J.J.,et al.Submitted (JAN-2000) to the EMBL/GenBank/DDBJ databases.
Bermudez-Fajardo A.,et al.Submitted (DEC-1999) to the EMBL/GenBank/DDBJ databases.
Humphray S.J.,et al.Nature 429:369-374(2004).