

Human Serum Albumin Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AP52692**Specification**

Human Serum Albumin Antibody - Product Information

Application	WB
Primary Accession	P02768
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	67 KDa

Human Serum Albumin Antibody - Additional Information**Gene ID** 213**Other Names**

ALB;ALBU_HUMAN;Albumin (32 AA);Albumin (AA 34);Albumin;Analbuminemia;Bisalbuminemia;Cell growth inhibiting protein 42;DKFZp779N1935;Dysalbuminemic hyperthyroxinemia;Growth inhibiting protein 20;HSA;Hyperthyroxinemia dysalbuminemic;PRO0883;PRO0903;PRO1341;PRO2044;PRO2619;Serum albumin.

Dilution

WB~~1:1000

Format

Purified mouse monoclonal in PBS(pH 7.4) containing with 0.09% (W/V) sodium azide and 50% glycerol.

Storage

Store at -20 °C.Stable for 12 months from date of receipt

Human Serum Albumin Antibody - Protein Information**Name** ALB**Function**

Binds water, Ca(2+), Na(+), K(+), fatty acids, hormones, bilirubin and drugs (Probable). Its main function is the regulation of the colloidal osmotic pressure of blood (Probable). Major zinc transporter in plasma, typically binds about 80% of all plasma zinc (PubMed:19021548). Major calcium and magnesium transporter in plasma, binds approximately 45% of circulating calcium and magnesium in plasma (By similarity). Potentially has more than two calcium-binding sites and might additionally bind calcium in a non-specific manner (By similarity). The shared binding site between zinc and calcium at residue Asp-273 suggests a crosstalk between zinc and calcium transport in the blood (By similarity). The rank order of affinity is zinc > calcium > magnesium (By

similarity). Binds to the bacterial siderophore enterobactin and inhibits enterobactin-mediated iron uptake of E.coli from ferric transferrin, and may thereby limit the utilization of iron and growth of enteric bacteria such as E.coli (PubMed:6234017). Does not prevent iron uptake by the bacterial siderophore aerobactin (PubMed:6234017).

Cellular Location

Secreted.

Tissue Location


Plasma.

Human Serum Albumin Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Human Serum Albumin Antibody - Images



Western blot detection of Human Serum Albumin in 0.5nl human serum and 5ng ALB Recombinant antigens cell lysates using Human Serum Albumin mouse mAb (1:1000 diluted). Predicted band size: 67KDa. Observed band size: 67KDa.

Human Serum Albumin Antibody - Background

Serum albumin, the main protein of plasma, has a good binding capacity for water, Ca(2+), Na(+), K(+), fatty acids, hormones, bilirubin and drugs. Its main function is the regulation of the colloidal osmotic pressure of blood. Major zinc transporter in plasma, typically binds about 80% of all plasma

zinc.

Human Serum Albumin Antibody - References

- Lawn R.M., et al. Nucleic Acids Res. 9:6103-6114(1981).
Dugaiczek A., et al. Proc. Natl. Acad. Sci. U.S.A. 79:71-75(1982).
Minghetti P.P., et al. J. Biol. Chem. 261:6747-6757(1986).
Yang S., et al. Submitted (SEP-1999) to the EMBL/GenBank/DDBJ databases.
Huang M.C., et al. Submitted (AUG-2002) to the EMBL/GenBank/DDBJ databases.