

Rad51 Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AP52714

Specification

Rad51 Antibody - Product Information

Application	WB
Primary Accession	Q06609
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a
Calculated MW	37 KDa

Rad51 Antibody - Additional Information

Gene ID 5888

Other Names

BRCA1/BRCA2 containing complex, subunit 5 ; BRCC 5 ; BRCC5 ; DNA repair protein RAD51 homolog 1 ; DNA repair protein rhp51 ; HRAD51 ; HsRad51 ; HsT16930 ; MRMV2 ; Rad 51 ; RAD51 ; RAD51 homolog (RecA homolog, E. coli) (S. cerevisiae) ; RAD51 homolog A ; RAD51 homolog ; RAD51 recombinase ; RAD51, S. cerevisiae, homolog of ; RAD51_HUMAN ; RAD51A ; RECA ; RecA like protein ; RecA, E. coli, homolog of ; Recombination protein A.

Dilution

WB~~1:1000

Format

0.09% (W/V) sodium azide and 50% glycerol.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Rad51 Antibody - Protein Information

Name RAD51 ([HGNC:9817](#))

Synonyms RAD51A, RECA

Function

Plays an important role in homologous strand exchange, a key step in DNA repair through homologous recombination (HR) (PubMed: [12205100](http://www.uniprot.org/citations/12205100) target="_blank">12205100, PubMed: [18417535](http://www.uniprot.org/citations/18417535) target="_blank">18417535, PubMed: [20231364](http://www.uniprot.org/citations/20231364) target="_blank">20231364, PubMed: [20348101](http://www.uniprot.org/citations/20348101) target="_blank">20348101, PubMed: [22325354](http://www.uniprot.org/citations/22325354) target="_blank">22325354, PubMed: [23509288](http://www.uniprot.org/citations/23509288) target="_blank">23509288)

target="_blank">23509288, PubMed:23754376, PubMed:26681308, PubMed:28575658, PubMed:32640219). Binds to single-stranded DNA in an ATP-dependent manner to form nucleoprotein filaments which are essential for the homology search and strand exchange (PubMed:12205100, PubMed:18417535, PubMed:20231364, PubMed:20348101, PubMed:23509288, PubMed:23754376, PubMed:26681308, PubMed:28575658). Catalyzes the recognition of homology and strand exchange between homologous DNA partners to form a joint molecule between a processed DNA break and the repair template (PubMed:12205100, PubMed:18417535, PubMed:20231364, PubMed:20348101, PubMed:23509288, PubMed:23754376, PubMed:26681308, PubMed:28575658, PubMed:38459011). Recruited to resolve stalled replication forks during replication stress (PubMed:27797818, PubMed:31844045). Part of a PALB2-scaffolded HR complex containing BRCA2 and RAD51C and which is thought to play a role in DNA repair by HR (PubMed:12442171, PubMed:24141787). Plays a role in regulating mitochondrial DNA copy number under conditions of oxidative stress in the presence of RAD51C and XRCC3 (PubMed:20413593). Also involved in interstrand cross-link repair (PubMed:26253028).

Cellular Location

Nucleus. Cytoplasm. Cytoplasm, perinuclear region. Mitochondrion matrix Chromosome. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome Note=Colocalizes with RAD51AP1 and RPA2 to multiple nuclear foci upon induction of DNA damage (PubMed:20154705). DNA damage induces an increase in nuclear levels (PubMed:20154705). Together with FIGNL1, redistributed in discrete nuclear DNA damage-induced foci after ionizing radiation (IR) or camptothecin (CPT) treatment (PubMed:23754376). Accumulated at sites of DNA damage in a SPIDR- dependent manner (PubMed:23509288). Recruited at sites of DNA damage in a MCM9-MCM8-dependent manner (PubMed:23401855). Recruited at sites of DNA damage following interaction with TOPBP1 in S-phase (PubMed:26811421). Colocalizes with ERCC5/XPG to nuclear foci in S phase (PubMed:26833090). Recruited to stalled replication forks during replication stress by the TONSL-MMS22L complex, as well as ATAD5 and WDR48 in an ATR-dependent manner (PubMed:27797818, PubMed:31844045)

Tissue Location


Highly expressed in testis and thymus, followed by small intestine, placenta, colon, pancreas and ovary. Weakly expressed in breast

Rad51 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Rad51 Antibody - Images



Western blot detection of Rad51 in CHO-K1, COS7, C6, 3T3 and HeLa cell lysates using Rad51 mouse mAb (1:1000 diluted). Predicted band size: 37KDa. Observed band size: 37KDa.

Rad51 Antibody - Background

Participates in a common DNA damage response pathway associated with the activation of homologous recombination and double-strand break repair. Binds to single and double-stranded DNA and exhibits DNA-dependent ATPase activity. Underwinds duplex DNA and forms helical nucleoprotein filaments. Part of a PALB2- scaffolded HR complex containing BRCA2 and RAD51C and which is thought to play a role in DNA repair by HR. Plays a role in regulating mitochondrial DNA copy number under conditions of oxidative stress in the presence of RAD51C and XRCC3.

Rad51 Antibody - References

- Shinohara A., et al. *Nat. Genet.* 4:239-243(1993).
Yoshimura Y., et al. *Nucleic Acids Res.* 21:1665-1665(1993).
Schmutte C., et al. *Cancer Res.* 59:4564-4569(1999).
Wang W.W., et al. *Cancer Epidemiol. Biomarkers Prev.* 10:955-960(2001).
Park J.Y., et al. *Nucleic Acids Res.* 36:3226-3234(2008).