

**CSK Antibody**  
**Purified Mouse Monoclonal Antibody (Mab)**  
**Catalog # AP52740**

**Specification**

---

**CSK Antibody - Product Information**

Application	<b>WB, ICC</b>
Primary Accession	<a href="#">P41240</a>
Reactivity	<b>Human</b>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>
Isotype	<b>IgG2b</b>
Calculated MW	<b>50 KDa</b>

**CSK Antibody - Additional Information**

**Gene ID** 1445

**Other Names**

C SRC;C SRC kinase;C src Tyrosine Kinase;C-SRC kinase;c-src tyrosine kinase;Csk A;CSK;  
CSK\_HUMAN;CYTOPLASMIC TYROSINE KINASE;EC  
2.7.10.2;MGC112926;MGC117393;MGC154049;P60 Src; Protein tyrosine kinase  
CYL;Protein-tyrosine kinase CYL;Proto oncogene tyrosine protein kinase;Tyrosine protein kinase  
CSK;Tyrosine protein kinase CSK;Tyrosine-protein kinase CSK;zgc:154049.

**Dilution**

WB~~1:1000  
ICC~~1:50

**Format**

Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine(pH 7.4,150 mM NaCl)with 0.09% (W/V) sodium azide,0.1mg/mlBSA and 50% glycerol.

**Storage**

Store at -20 °C.Stable for 12 months from date of receipt

**CSK Antibody - Protein Information**

**Name** CSK

**Function**

Non-receptor tyrosine-protein kinase that plays an important role in the regulation of cell growth, differentiation, migration and immune response. Phosphorylates tyrosine residues located in the C-terminal tails of Src-family kinases (SFKs) including LCK, SRC, HCK, FYN, LYN, CSK or YES1. Upon tail phosphorylation, Src-family members engage in intramolecular interactions between the phosphotyrosine tail and the SH2 domain that result in an inactive conformation. To inhibit SFKs, CSK is recruited to the plasma membrane via binding to transmembrane proteins or adapter proteins located near the plasma membrane. Suppresses signaling by various surface receptors,

including T-cell receptor (TCR) and B-cell receptor (BCR) by phosphorylating and maintaining inactive several positive effectors such as FYN or LCK.

#### Cellular Location

Cytoplasm. Cell membrane. Note=Mainly cytoplasmic, also present in lipid rafts

#### Tissue Location


Expressed in lung and macrophages.

### CSK Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CSK Antibody - Images



Western blot detection of CSK in C6 and Jurkat cell lysates using CSK mouse mAb (1:1000 diluted). Predicted band size: 50KDa. Observed band size: 50KDa.



Immunocytochemistry staining of C6 cells fixed by anhydrous methanol for 2 h at -20°C and using anti-CSK mouse mAb (dilution 1:50).

### **CSK Antibody - Background**

Non-receptor tyrosine-protein kinase that plays an important role in the regulation of cell growth, differentiation, migration and immune response. Phosphorylates tyrosine residues located in the C-terminal tails of Src-family kinases (SFks) including LCK, SRC, HCK, FYN, LYN or YES1. Upon tail phosphorylation, Src-family members engage in intramolecular interactions between the phosphotyrosine tail and the SH2 domain that result in an inactive conformation. To inhibit SFks, CSK is recruited to the plasma membrane via binding to transmembrane proteins or adapter proteins located near the plasma membrane. Suppresses signaling by various surface receptors, including T- cell receptor (TCR) and B-cell receptor (BCR) by phosphorylating and maintaining inactive several positive effectors such as FYN or LCK.

### **CSK Antibody - References**

Partanen J., et al. *Oncogene* 6:2013-2018(1991).  
Braeuninger A., et al. *Proc. Natl. Acad. Sci. U.S.A.* 88:10411-10415(1991).  
Brauninger A., et al. *Gene* 110:205-211(1992).  
Braeuninger A., et al. *Oncogene* 8:1365-1369(1993).  
Halleck A., et al. Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.