

**FEN-1 Antibody**  
**Purified Mouse Monoclonal Antibody (Mab)**  
**Catalog # AP52835****Specification**

---

**FEN-1 Antibody - Product Information**

Application	<b>WB, ICC</b>
Primary Accession	<a href="#">P39748</a>
Reactivity	<b>Human, Mouse</b>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>
Isotype	<b>IgG1</b>
Calculated MW	<b>45 KDa</b>

**FEN-1 Antibody - Additional Information****Gene ID** 2237**Other Names**

DNase IV;FEN-1;FEN1;FEN1\_HUMAN;Flap endonuclease 1;Flap structure specific endonuclease 1;Flap structure-specific endonuclease 1;hFEN-1;hFEN1;Maturation factor 1;MF1;Rad2.

**Dilution**

WB~~1:1000

ICC~~1:400

**Format**

Purified mouse monoclonal antibody in PBS(pH 7.4) containing with 0.09% (W/V) sodium azide and 50% glycerol.

**Storage**

Store at -20 °C.Stable for 12 months from date of receipt

**FEN-1 Antibody - Protein Information****Name** FEN1 {ECO:0000255|HAMAP-Rule:MF\_03140}**Synonyms** RAD2**Function**

Structure-specific nuclease with 5'-flap endonuclease and 5'- 3' exonuclease activities involved in DNA replication and repair. During DNA replication, cleaves the 5'-overhanging flap structure that is generated by displacement synthesis when DNA polymerase encounters the 5'-end of a downstream Okazaki fragment. It enters the flap from the 5'-end and then tracks to cleave the flap base, leaving a nick for ligation. Also involved in the long patch base excision repair (LP-BER) pathway, by cleaving within the apurinic/aprimidinic (AP) site- terminated flap. Acts as a genome stabilization factor that prevents flaps from equilibrating into structures that lead to duplications and deletions. Also possesses 5'-3' exonuclease activity on nicked or gapped double-stranded

DNA, and exhibits RNase H activity. Also involved in replication and repair of rDNA and in repairing mitochondrial DNA.

#### Cellular Location


[Isoform 1]: Nucleus, nucleolus. Nucleus, nucleoplasm. Note=Resides mostly in the nucleoli and relocalizes to the nucleoplasm upon DNA damage

#### FEN-1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### FEN-1 Antibody - Images



Western blot detection of FEN-1 in HeLa, Jurkat, 3T3, COS7, PC-12, C6, Raw264.7, Raji and CHO-K1 cell lysates using FEN-1 mouse mAb (1:1000 diluted). Predicted band size: 45 kDa. Observed band size: 45 kDa.



Immunocytochemistry staining of HeLa cells fixed with 4% Paraformaldehyde and using FEN-1 mouse mAb (dilution 1:400).

### **FEN-1 Antibody - Background**

Structure-specific nuclease with 5'-flap endonuclease and 5'-3' exonuclease activities involved in DNA replication and repair. During DNA replication, cleaves the 5'-overhanging flap structure that is generated by displacement synthesis when DNA polymerase encounters the 5'-end of a downstream Okazaki fragment. It enters the flap from the 5'-end and then tracks to cleave the flap base, leaving a nick for ligation. Also involved in the long patch base excision repair (LP-BER) pathway, by cleaving within the apurinic/aprimidinic (AP) site-terminated flap. Acts as a genome stabilization factor that prevents flaps from equilibrating into structures that lead to duplications and deletions. Also possesses 5'-3' exonuclease activity on nicked or gapped double-stranded DNA, and exhibits RNase H activity. Also involved in replication and repair of rDNA and in repairing mitochondrial DNA.

### **FEN-1 Antibody - References**

Murray J.M., et al. Mol. Cell. Biol. 14:4878-4888(1994).  
Hiraoka L.R., et al. Genomics 25:220-225(1995).  
Taylor T.D., et al. Nature 440:497-500(2006).  
Robins P., et al. J. Biol. Chem. 269:28535-28538(1994).  
Shen B., et al. J. Biol. Chem. 271:9173-9176(1996).