

**SAR1B Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP53321****Specification**

---

**SAR1B Antibody - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">O9Y6B6</a>
Reactivity	<b>Human, Mouse</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>22 KDa</b>
Antigen Region	<b>116-165</b>

**SAR1B Antibody - Additional Information****Gene ID** 51128**Other Names**

GTP-binding protein SAR1b, GTP-binding protein B, GTBPB, SAR1B, SARA2, SARB

**Dilution**

WB~~ 1:1000

**Format**

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**SAR1B Antibody - Protein Information****Name** SAR1B {ECO:0000303|PubMed:33186557, ECO:0000312|HGNC:HGNC:10535}**Function**

Small GTPase that cycles between an active GTP-bound and an inactive GDP-bound state and mainly functions in vesicle-mediated endoplasmic reticulum (ER) to Golgi transport. The active GTP-bound form inserts into the endoplasmic reticulum membrane where it recruits the remainder of the coat protein complex II/COPII (PubMed: [23433038](http://www.uniprot.org/citations/23433038), PubMed: [32358066](http://www.uniprot.org/citations/32358066), PubMed: [33186557](http://www.uniprot.org/citations/33186557), PubMed: [36369712](http://www.uniprot.org/citations/36369712)). The coat protein complex II assembling and polymerizing on endoplasmic reticulum membrane is responsible for both the sorting of cargos and the deformation and budding of membranes into vesicles destined to the Golgi (PubMed: [23433038](http://www.uniprot.org/citations/23433038), PubMed: [32358066](http://www.uniprot.org/citations/32358066), PubMed: [33186557](http://www.uniprot.org/citations/33186557), PubMed: [36369712](http://www.uniprot.org/citations/36369712)).

href="http://www.uniprot.org/citations/33186557" target="\_blank">33186557</a>). In contrast to SAR1A, SAR1B specifically interacts with the cargo receptor SURF4 to mediate the transport of lipid-carrying lipoproteins including APOB and APOA1 from the endoplasmic reticulum to the Golgi and thereby, indirectly regulates lipid homeostasis (PubMed:<a href="http://www.uniprot.org/citations/32358066" target="\_blank">32358066</a>, PubMed:<a href="http://www.uniprot.org/citations/33186557" target="\_blank">33186557</a>). In addition to its role in vesicle trafficking, can also function as a leucine sensor regulating TORC1 signaling and more indirectly cellular metabolism, growth and survival. In absence of leucine, interacts with the GATOR2 complex via MIOS and inhibits TORC1 signaling. The binding of leucine abrogates the interaction with GATOR2 and the inhibition of the TORC1 signaling. This function is completely independent of the GTPase activity of SAR1B (PubMed:<a href="http://www.uniprot.org/citations/34290409" target="\_blank">34290409</a>).

#### Cellular Location

Endoplasmic reticulum membrane; Peripheral membrane protein {ECO:0000250|UniProtKB:Q9QVY3}. Golgi apparatus, Golgi stack membrane {ECO:0000250|UniProtKB:Q9QVY3}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q9QVY3}. Cytoplasm, cytosol. Lysosome membrane. Note=Active at endoplasmic reticulum exit sites (ERES) where it inserts into the membrane and recruits the remainder of the coat protein complex II/COPII (PubMed:23433038). Upon leucine deprivation, associates with lysosomal membranes to repress TORC1 signaling (PubMed:34290409).

#### Tissue Location

Expressed in many tissues including small intestine, liver, muscle and brain.


#### SAR1B Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### SAR1B Antibody - Images





Anti-SAR1B Antibody at 1:1000 dilution + HeLa whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 22 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

### **SAR1B Antibody - Background**

Involved in transport from the endoplasmic reticulum to the Golgi apparatus. Activated by the guanine nucleotide exchange factor PREB. Involved in the selection of the protein cargo and the assembly of the COPII coat complex.

### **SAR1B Antibody - References**

Song H., et al. Submitted (SEP-1998) to the EMBL/GenBank/DDBJ databases.  
Zhou Y., et al. Submitted (JUL-2003) to the EMBL/GenBank/DDBJ databases.  
Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.  
Burkard T.R., et al. BMC Syst. Biol. 5:17-17(2011).  
Jones B., et al. Nat. Genet. 34:29-31(2003).