

Anti-EpCAM Antibody
Mouse Anti Human Monoclonal Antibody
Catalog # AP53386

Specification

Anti-EpCAM Antibody - Product Information

Application	IP, WB
Primary Accession	P16422
Other Accession	NM_002354
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Immunogen	Purified recombinant EpCAM protein fragments expressed in E.coli.
Purification	Affinity purified
Calculated MW	39 KDa

Anti-EpCAM Antibody - Additional Information

Gene ID 4072

Other Names

17 1A; 323/A3; Adenocarcinoma associated antigen; Adenocarcinoma-associated antigen; Antigen identified by monoclonal antibody AUA1; AUA1; CD326; CD326 antigen; Cell surface glycoprotein Trop 1; Cell surface glycoprotein Trop 2; Cell surface glycoprotein Trop-1; CO 17A; CO17 1A; CO17A; DIAR5; EGP 2; EGP; EGP2; EGP314; EGP40; Ep CAM; Ep-CAM; EPCAM; EPCAM_HUMAN; Epithelial cell adhesion molecule; Epithelial Cell Adhesion Molecule Intracellular Domain (EpCAM-ICD); Epithelial cell surface antigen; Epithelial cellular adhesion molecule; Epithelial glycoprotein 1; Epithelial glycoprotein 314; Epithelial glycoprotein; ESA; GA733 1; GA733 2; GA733-2; gastrointestinal tumor-associated antigen 2, 35-KD glycoprotein; hEGP 2; hEGP314; HNPCC8; Human epithelial glycoprotein 2; KS 1/4 antigen; KS1/4; KSA; Lymphocyte antigen 74; M1S 1; M1S2; M4S1; Major gastrointestinal tumor associated protein GA733 2; Major gastrointestinal tumor-associated protein GA733-2; Membrane component chromosome 4 surface marker (35kD glycoprotein); Membrane component, chromosome 4, surface marker 1; Membrane component, chromosome 4, surface marker; MIC18; MK 1; TACD1; TACSTD1; TROP1; Tumor associated calcium signal transducer 1; Tumor associated calcium signal transducer 2 precursor; Tumor-associated calcium signal transducer 1.

Dilution

WB~~1:1000

Format

PBS(pH 7.4) containing with 0.09% (W/V) sodium azide and 50% glycerol.

Storage

Store at -20 °C.Stable for 12 months from date of receipt

Anti-EpCAM Antibody - Protein Information

Name EPCAM

Synonyms GA733-2, M1S2, M4S1, MIC18, TACSTD1, TRO

Function

May act as a physical homophilic interaction molecule between intestinal epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium for providing immunological barrier as a first line of defense against mucosal infection. Plays a role in embryonic stem cells proliferation and differentiation. Up-regulates the expression of FABP5, MYC and cyclins A and E.

Cellular Location

Lateral cell membrane; Single-pass type I membrane protein. Cell junction, tight junction.
Note=Colocalizes with CLDN7 at the lateral cell membrane and tight junction

Tissue Location


Highly and selectively expressed by undifferentiated rather than differentiated embryonic stem cells (ESC) Levels rapidly diminish as soon as ESC's differentiate (at protein levels). Expressed in almost all epithelial cell membranes but not on mesodermal or neural cell membranes. Found on the surface of adenocarcinoma.

Anti-EpCAM Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-EpCAM Antibody - Images



Immunoprecipitation analysis of HCT116 and HeLa cell lysates using EpCAM.



Western blot analysis of extracts from CHO-K1, CHO-K1 transfected by Flag-EpCAM, MCF7 (EpCAM positive), HCT116 (EpCAM positive), and HeLa (EpCAM negative) cell lysates using EpCAM mouse mAb (1:1000 diluted). Predicted band size: 39 kDa. Observed band size: 39/35 kDa.

Anti-EpCAM Antibody - Background

May act as a physical homophilic interaction molecule between intestinal epithelial cells (IECs) and intraepithelial lymphocytes (IELs) at the mucosal epithelium for providing immunological barrier as a first line of defense against mucosal infection. Pla