10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Anti-MMP-2 Antibody
Mouse Anti Human Monoclonal Antibody
Catalog # AP53416

Specification

Anti-MMP-2 Antibody - Product Information

Application	**WB, ICC**
Primary Accession	P08253
Other Accession	NM_004530
Reactivity	**Human, Rat**
Host	**Mouse**
Clonality	**Monoclonal**
Isotype	**IgG1**
Immunogen	**Purified recombinant human MMP-2 protein expressed in E.coli.**
Purification	**Affinity purified**
Calculated MW	**72 KDa**

Anti-MMP-2 Antibody - Additional Information

Gene ID 4313

Other Names
72 kDa gelatinase;72kD type IV collagenase;CLG 4;CLG 4A;CLG4;CLG4A;Collagenase Type 4 alpha;Collagenase type IV A;Gelatinase A;Gelatinase alpha;Gelatinase neutrophil;Matrix metallopeptidase 2 gelatinase A 72kDa gelatinase 72kDa type IV collagenase;Matrix metalloproteinase 2 (gelatinase A, 72kDa gelatinase, 72kDa type IV collagenase);Matrix Metalloproteinase 2;Matrix metalloproteinase II;Matrix metalloproteinase-2;MMP 2;MMP II;MMP-2;MMP2;MMP2_HUMAN;MONA;Neutrophil gelatinase;PEX;TBE 1;TBE-1.

Dilution
WB~~1:1000

Format
PBS(pH 7.4) containing with 0.09% (W/V) sodium azide and 50% glycerol.

Storage
Store at -20 °C.Stable for 12 months from date of receipt

Anti-MMP-2 Antibody - Protein Information

Name MMP2

Synonyms CLG4A

Function
Ubiquitinous metalloproteinase that is involved in diverse functions such as remodeling of the vasculature, angiogenesis, tissue repair, tumor invasion, inflammation, and atherosclerotic plaque

rupture. As well as degrading extracellular matrix proteins, can also act on several nonmatrix proteins such as big endothelial 1 and beta- type CGRP promoting vasoconstriction. Also cleaves KISS at a Gly-|-Leu bond. Appears to have a role in myocardial cell death pathways. Contributes to myocardial oxidative stress by regulating the activity of GSK3beta. Cleaves GSK3beta in vitro. Involved in the formation of the fibrovascular tissues in association with MMP14. [Isoform 2]: Mediates the proteolysis of CHUK/IKKA and initiates a primary innate immune response by inducing mitochondrial- nuclear stress signaling with activation of the pro-inflammatory NF-kappaB, NFAT and IRF transcriptional pathways.

Cellular Location
[Isoform 1]: Secreted, extracellular space, extracellular matrix. Membrane. Nucleus Note=Colocalizes with integrin alphaV/beta3 at the membrane surface in angiogenic blood vessels and melanomas. Found in mitochondria, along microfibrils, and in nuclei of cardiomyocytes

Tissue Location
Produced by normal skin fibroblasts. PEX is expressed in a number of tumors including gliomas, breast and prostate

Anti-MMP-2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-MMP-2 Antibody - Images



Western blot analysis of extracts from CHO-K1 (B) and CHO-K1 transfected by MMP-2 fragment(A) cell lysates using MMP-2 mouse mAb (1:2000 diluted). Predicted band size:30KDa. Observed band size:30KDa.

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



Western blot analysis of extracts from Raw264.7 cell lysates using MMP-2 mouse mAb (1:200 diluted). Predicted band size:64,72KDa. Observed band size:72KDa.



Immunofluorescent analysis of Hela cells fixed fixed by anhydrous methanol at -20℃ and using MMP-2 mouse mAb (dilution 1:50). DAPI was used to stain nucleus (blue).

Anti-MMP-2 Antibody - Background

Ubiquitinous metalloproteinase that is involved in diverse functions such as remodeling of the vasculature, angiogenesis, tissue repair, tumor invasion, inflammation, and atherosclerotic plaque rupture. As well as degrading extracellular matrix proteins,

Anti-MMP-2 Antibody - Citations

- Circ_0046599 Promotes the Development of Hepatocellular Carcinoma by Regulating the miR-1258/RPN2 Network