

HIST1H2AK Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5347A

Specification

HIST1H2AK Antibody (N-term) - Product Information

| | |
|-------------------|--|
| Application | WB, E |
| Primary Accession | P0C0S8 |
| Other Accession | P02263 , Q4FZT6 , Q8BFU2 , Q7L7L0 , P35062 , Q64523 , Q16777 , A1A4R1 , Q64522 , Q8IUE6 , P0CC09 , Q6GSS7 , Q6FI13 , P02262 , P22752 , P0C0S9 , Q8CGP7 , Q99878 , Q8CGP6 , Q96KK5 , Q64598 , Q8CGP5 , P0C170 , P20671 , P0C169 , Q93077 , P04908 , NP_003501.1 , C0HKE1 , C0HKE2 , C0HKE3 , C0 |
| Reactivity | Human |
| Predicted | Mouse, Rat, Bovine, Chicken |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Calculated MW | 14091 |
| Antigen Region | 1-30 |

HIST1H2AK Antibody (N-term) - Additional Information

Gene ID 8329;8330;8332;8336;8969

Other Names

Histone H2A type 1, H2A1, Histone H2A/p, HIST1H2AG, H2AFP

Target/Specificity

This HIST1H2AK antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human HIST1H2AK.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HIST1H2AK Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

HIST1H2AK Antibody (N-term) - Protein Information

Name H2AC11 ([HGNC:4737](#))

Synonyms H2AFP, HIST1H2AG

Function Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Cellular Location


Nucleus. Chromosome.

HIST1H2AK Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HIST1H2AK Antibody (N-term) - Images



HIST1H2AK Antibody (N-term) (Cat. #AP5347a) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the HIST1H2AK antibody detected the HIST1H2AK protein (arrow).

HIST1H2AK Antibody (N-term) - Background

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless

and encodes a member of the histone H2A family.

HIST1H2AK Antibody (N-term) - References

Bergink, S., et al. Genes Dev. 20(10):1343-1352(2006)
Bonenfant, D., et al. Mol. Cell Proteomics 5(3):541-552(2006)
Boyne, M.T. II, et al. J. Proteome Res. 5(2):248-253(2006)