

HMGA2 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5359b**Specification**

HMGA2 Antibody (C-term) - Product Information

Application	WB, FC,E
Primary Accession	P52926
Other Accession	P52927 , NP_003475.1 , NP_003474.1
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	11832
Antigen Region	64-92

HMGA2 Antibody (C-term) - Additional Information**Gene ID** 8091**Other Names**

High mobility group protein HMGI-C, High mobility group AT-hook protein 2, HMGA2, HMGIC

Target/Specificity

This HMGA2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 64-92 amino acids from the C-terminal region of human HMGA2.

DilutionWB~~1:1000
FC~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HMGA2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

HMGA2 Antibody (C-term) - Protein Information**Name** HMGA2

Synonyms HMGIC

Function Functions as a transcriptional regulator. Functions in cell cycle regulation through CCNA2. Plays an important role in chromosome condensation during the meiotic G2/M transition of spermatocytes. Plays a role in postnatal myogenesis, is involved in satellite cell activation (By similarity). Positively regulates IGF2 expression through PLAG1 and in a PLAG1-independent manner (PubMed:[28796236](#)).

Cellular Location


Nucleus.

HMGA2 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HMGA2 Antibody (C-term) - Images



HMGA2 Antibody (C-term) (Cat. #AP5359b) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the HMGA2 antibody detected the HMGA2 protein (arrow).



HMGA2 Antibody (C-term) (Cat. #AP5359b) flow cytometric analysis of HepG2 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

HMGA2 Antibody (C-term) - Background

This gene encodes a protein that belongs to the non-histone chromosomal high mobility group (HMG) protein family. HMG proteins function as architectural factors and are essential components of the enhancosome. This protein contains structural DNA-binding domains and may act as a transcriptional regulating factor. Identification of the deletion, amplification, and rearrangement of this gene that are associated with myxoid liposarcoma suggests a role in adipogenesis and mesenchymal differentiation. A gene knock out study of the mouse counterpart demonstrated that this gene is involved in diet-induced obesity. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.

HMGA2 Antibody (C-term) - References

Mu, G., et al. Hum. Pathol. 41(4):493-502(2010)
Pillas, D., et al. PLoS Genet. 6 (2), E1000856 (2010) :
Yang, T.L., et al. Ann. Hum. Genet. 74(1):11-16(2010)
Wei, J.J., et al. Am. J. Surg. Pathol. 34(1):18-26(2010)
Tay, Y., et al. Stem Cell Rev 5(4):328-333(2009)
Schwanbeck, R., et al. J. Biol. Chem. 275(3):1793-1801(2000)
Chau, K.Y., et al. Nucleic Acids Res. 23(21):4262-4266(1995)
Schoenmakers, E.F., et al. Nat. Genet. 10(4):436-444(1995)
Ashar, H.R., et al. Cell 82(1):57-65(1995)
Schoenmakers, E.F., et al. Genes Chromosomes Cancer 11(2):106-118(1994)
Manfioletti, G., et al. Nucleic Acids Res. 19(24):6793-6797(1991)

HMGA2 Antibody (C-term) - Citations

- [Cr \(VI\) induced mitophagy via the interaction of HMGA2 and PARK2](#)
- [HMGA2 upregulation mediates Cd-induced migration and invasion in A549 cells and in lung tissues of mice.](#)