

**Anti-ACTN2 Antibody**  
Catalog # AP53654

**Specification**

---

**Anti-ACTN2 Antibody - Product Information**

Application	<b>WB, IF</b>
Primary Accession	<a href="#">P35609</a>
Other Accession	<a href="#">Q08043</a>
Reactivity	<b>Human, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>103854</b>

**Anti-ACTN2 Antibody - Additional Information**

**Gene ID** 88

**Other Names**

ACTN2; Alpha-actinin-2; Alpha-actinin skeletal muscle isoform 2; F-actin cross-linking protein; ACTN3; Alpha-actinin-3; Alpha-actinin skeletal muscle isoform 3; F-actin cross-linking protein

**Target/Specificity**

Recognizes endogenous levels of ACTN2 protein.

**Dilution**

WB~~1/500 - 1/1000  
IF~~1/50 - 1/200

**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-ACTN2 Antibody - Protein Information**

**Name** ACTN2

**Function**

F-actin cross-linking protein which is thought to anchor actin to a variety of intracellular structures. This is a bundling protein.

**Cellular Location**

Cytoplasm, myofibril, sarcomere, Z line. Note=Colocalizes with MYOZ1 and FLNC at the Z-lines of skeletal muscle

**Tissue Location**


Expressed in both skeletal and cardiac muscle.

### Anti-ACTN2 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### Anti-ACTN2 Antibody - Images



Western blot analysis of ACTN2 expression in HeLa (A), mouse kidney (B), H9C2 (C) whole cell lysates.



Immunohistochemical analysis of ACTN2 staining in human muscle formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of ACTN2 staining in H9C2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

#### **Anti-ACTN2 Antibody - Background**

Rabbit polyclonal antibody to ACTN2