

**Anti-p67 Antibody**  
Catalog # AP53900**Specification**

---

**Anti-p67 Antibody - Product Information**

Application	WB, IF
Primary Accession	<a href="#">P61764</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67569

**Anti-p67 Antibody - Additional Information****Gene ID** 6812**Other Names**

UNC18A; Syntaxin-binding protein 1; MUNC18-1; N-Sec1; Protein unc-18 homolog 1; Unc18-1; Protein unc-18 homolog A; Unc-18A; p67

**Target/Specificity**

Recognizes endogenous levels of p67 protein.

**Dilution**WB~~1/500 - 1/1000  
IF~~1/50 - 1/200**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-p67 Antibody - Protein Information****Name** STXBP1**Synonyms** UNC18A**Function**

Participates in the regulation of synaptic vesicle docking and fusion through interaction with GTP-binding proteins (By similarity). Essential for neurotransmission and binds syntaxin, a component of the synaptic vesicle fusion machinery probably in a 1:1 ratio. Can interact with syntaxins 1, 2, and 3 but not syntaxin 4. Involved in the release of neurotransmitters from neurons through interacting with SNARE complex component STX1A and mediating the assembly of the SNARE complex at synaptic membranes (By similarity). May play a role in determining the specificity of intracellular fusion reactions.

**Cellular Location**

Cytoplasm, cytosol. Membrane; Peripheral membrane protein


**Tissue Location**

Brain and spinal cord. Highly enriched in axons.


**Anti-p67 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


**Anti-p67 Antibody - Images**

Western blot analysis of p67 expression in U87MG (A), C6 (B), SHSY5Y (C), K562 (D), Raw264.7 (E), mouse brain (F), rat brain (G) whole cell lysates.



Immunohistochemical analysis of p67 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium

citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of p67 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

#### **Anti-p67 Antibody - Background**

Rabbit polyclonal antibody to p67