

Anti-Glycerate Kinase Antibody
Catalog # AP53912**Specification**

Anti-Glycerate Kinase Antibody - Product Information

Application	WB, IF
Primary Accession	Q8IVS8
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55253

Anti-Glycerate Kinase Antibody - Additional Information**Gene ID** 132158**Other Names**

HBEBP4; Glycerate kinase; HBeAg-binding protein 4

Target/Specificity

Recognizes endogenous levels of Glycerate Kinase protein.

Dilution

WB~~1/500 - 1/1000

IF~~1/50 - 1/200

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-Glycerate Kinase Antibody - Protein Information**Name** GLYCTK**Synonyms** HBEBP4**Cellular Location**

[Isoform 1]: Cytoplasm

Tissue Location


Widely expressed..

Anti-Glycerate Kinase Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-Glycerate Kinase Antibody - Images



Western blot analysis of Glycerate Kinase expression in HepG2 (A), NIH3T3 (B) whole cell lysates.



Immunohistochemical analysis of Glycerate Kinase staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Glycerate Kinase staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Anti-Glycerate Kinase Antibody - Background

Rabbit polyclonal antibody to Glycerate Kinase