

**Anti-PEX19 Antibody**  
Catalog # AP53920

**Specification**

---

**Anti-PEX19 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P40855</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32807

**Anti-PEX19 Antibody - Additional Information**

Gene ID 5824

**Other Names**

HK33; PXF; Peroxisomal biogenesis factor 19; 33 kDa housekeeping protein; Peroxin-19; Peroxisomal farnesylated protein

**Target/Specificity**

Recognizes endogenous levels of PEX19 protein.

**Dilution**

WB~~1/500 - 1/1000

**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-PEX19 Antibody - Protein Information**

Name PEX19 ([HGNC:9713](#))

Synonyms HK33, PXF

**Function**

Necessary for early peroxisomal biogenesis. Acts both as a cytosolic chaperone and as an import receptor for peroxisomal membrane proteins (PMPs). Binds and stabilizes newly synthesized PMPs in the cytoplasm by interacting with their hydrophobic membrane-spanning domains, and targets them to the peroxisome membrane by binding to the integral membrane protein PEX3. Excludes CDKN2A from the nucleus and prevents its interaction with MDM2, which results in active degradation of TP53.

**Cellular Location**

Cytoplasm. Peroxisome membrane; Lipid-anchor; Cytoplasmic side. Note=Mainly cytoplasmic. Some fraction membrane-associated to the outer surface of peroxisomes.

#### Tissue Location


Ubiquitously expressed. Isoform 1 is strongly predominant in all tissues except in utero where isoform 2 is the main form.

#### Anti-PEX19 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Anti-PEX19 Antibody - Images



Western blot analysis of PEX19 expression in HEK293T (A), mouse brain (B), rat brain (C) whole cell lysates.



Immunohistochemical analysis of PEX19 staining in human lung cancer formalin fixed paraffin

embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

#### **Anti-PEX19 Antibody - Background**

Rabbit polyclonal antibody to PEX19