

Anti-ZNF392 Antibody
Catalog # AP53950**Specification**

Anti-ZNF392 Antibody - Product Information

Application	WB, IF
Primary Accession	Q9H4T2
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40792

Anti-ZNF392 Antibody - Additional Information**Gene ID** 80345**Other Names**ZNF392; ZNF435; Zinc finger and SCAN domain-containing protein 16; Zinc finger protein 392;
Zinc finger protein 435**Target/Specificity**

Recognizes endogenous levels of ZNF392 protein.

DilutionWB~~1/500 - 1/1000
IF~~1/50 - 1/200**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-ZNF392 Antibody - Protein Information**Name** ZSCAN16**Synonyms** ZNF392, ZNF435**Function**

May be involved in transcriptional regulation.

Cellular Location


Nucleus.

Anti-ZNF392 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-ZNF392 Antibody - Images



Western blot analysis of ZNF392 expression in HCT116 (A), DLD (B), mouse brain (C) whole cell lysates.



Immunohistochemical analysis of ZNF392 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of ZNF392 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with Alexa Fluor 647-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Anti-ZNF392 Antibody - Background

Rabbit polyclonal antibody to ZNF392