

**Anti-NANOG Antibody**  
Catalog # AP54003**Specification****Anti-NANOG Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q9H9S0</a>
Other Accession	<a href="#">Q8N7R0</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34620

**Anti-NANOG Antibody - Additional Information****Gene ID** 79923**Other Names**

Homeobox protein NANOG; Homeobox transcription factor Nanog; hNanog

**Target/Specificity**

Recognizes endogenous levels of NANOG protein.

**Dilution**

WB~~1/500 - 1/1000

**Format**

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Anti-NANOG Antibody - Protein Information****Name** NANOG**Function**

Transcription regulator involved in inner cell mass and embryonic stem (ES) cells proliferation and self-renewal. Imposes pluripotency on ES cells and prevents their differentiation towards extraembryonic endoderm and trophoctoderm lineages. Blocks bone morphogenetic protein-induced mesoderm differentiation of ES cells by physically interacting with SMAD1 and interfering with the recruitment of coactivators to the active SMAD transcriptional complexes. Acts as a transcriptional activator or repressor. Binds optimally to the DNA consensus sequence 5'-TAAT[GT][GT]-3' or 5'-[CG][GA][CG]C[GC]ATTAN[GC]- 3'. Binds to the POU5F1/OCT4 promoter (PubMed:<a href="http://www.uniprot.org/citations/25825768" target="\_blank">25825768</a>). Able to autorepress its expression in differentiating (ES) cells: binds to its own promoter following interaction with ZNF281/ZFP281, leading to recruitment of the NuRD complex and subsequent

repression of expression. When overexpressed, promotes cells to enter into S phase and proliferation.

#### Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108, ECO:0000269|PubMed:15983365}

#### Tissue Location


Expressed in testicular carcinoma and derived germ cell tumors (at protein level). Expressed in fetal gonads, ovary and testis. Also expressed in ovary teratocarcinoma cell line and testicular embryonic carcinoma. Not expressed in many somatic organs and oocytes.

### Anti-NANOG Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-NANOG Antibody - Images



Western blot analysis of NANOG expression in mouse embryo (A), rat testis (B), A2780 (C), HeLa (D) whole cell lysates.

### Anti-NANOG Antibody - Background

Rabbit polyclonal antibody to NANOG