

Anti-TAF4 Antibody
Catalog # AP54061**Specification**

Anti-TAF4 Antibody - Product Information

Application	WB
Primary Accession	O00268
Other Accession	Q92750
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	110114

Anti-TAF4 Antibody - Additional Information**Gene ID** 6874**Other Names**

TAF4; TAF2C; TAF2C1; TAF4A; TAFII130; TAFII135; Transcription initiation factor TFIID subunit 4; RNA polymerase II TBP-associated factor subunit C; TBP-associated factor 4; Transcription initiation factor TFIID 130 kDa subunit; TAF(II)130; TAFII-130; TAFII130; Transcription initiation factor TFIID 135 kDa subunit; TAF(II)135; TAFII-135; TAFII135; TAF4B; TAF2C2; TAFII105; Transcription initiation factor TFIID subunit 4B; Transcription initiation factor TFIID 105 kDa subunit; TAF(II)105; TAFII-105; TAFII105

Target/Specificity

Recognizes endogenous levels of TAF4 protein.

Dilution

WB~~1/500 - 1/1000

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Anti-TAF4 Antibody - Protein Information**Name** TAF4**Synonyms** TAF2C, TAF2C1, TAF4A, TAFII130, TAFII135**Function**

The TFIID basal transcription factor complex plays a major role in the initiation of RNA polymerase II (Pol II)-dependent transcription (PubMed: 33795473). TFIID recognizes and binds promoters with or without a TATA

box via its subunit TBP, a TATA-box-binding protein, and promotes assembly of the pre-initiation complex (PIC) (PubMed:33795473). The TFIID complex consists of TBP and TBP-associated factors (TAFs), including TAF1, TAF2, TAF3, TAF4, TAF5, TAF6, TAF7, TAF8, TAF9, TAF10, TAF11, TAF12 and TAF13 (PubMed:10594036, PubMed:33795473, PubMed:8942982). TAF4 may maintain an association between the TFIID and TFIIA complexes, while bound to the promoter, together with TBP, during PIC assembly (PubMed:33795473). Potentiates transcriptional activation by the AF-2S of the retinoic acid, vitamin D3 and thyroid hormone (PubMed:9192867).

Cellular Location


Nucleus.

Anti-TAF4 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-TAF4 Antibody - Images



Western blot analysis of TAF4 expression in A2780 (A), SKOV3 (B), mouse embryo (C) whole cell lysates.



Immunohistochemical analysis of TAF4 staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-TAF4 Antibody - Background

Rabbit polyclonal antibody to TAF4