

Neurokinin A Receptor Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54183**Specification**

Neurokinin A Receptor Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	P21452
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	44442

Neurokinin A Receptor Polyclonal Antibody - Additional Information**Gene ID** 6865**Other Names**

Substance-K receptor, SKR, NK-2 receptor, NK-2R, Neurokinin A receptor, Tachykinin receptor 2, TACR2, NK2R, NKNAR, TAC2R

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Neurokinin A Receptor Polyclonal Antibody - Protein Information**Name** TACR2**Synonyms** NK2R, NKNAR, TAC2R**Function**

This is a receptor for the tachykinin neuropeptide substance K (neurokinin A). It is associated with G proteins that activate a phosphatidylinositol-calcium second messenger system. The rank order of affinity of this receptor to tachykinins is: substance K > neuromedin-K > substance P.

Cellular Location


Cell membrane; Multi-pass membrane protein.

Neurokinin A Receptor Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Neurokinin A Receptor Polyclonal Antibody - Images



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-NK-2R Polyclonal Antibody, Unconjugated(bs-0123R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining