

PTOV1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54231

Specification

PTOV1 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q86YD1
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	46869

PTOV1 Polyclonal Antibody - Additional Information

Gene ID 53635

Other Names

Prostate tumor-overexpressed gene 1 protein, PTOV-1, Activator interaction domain-containing protein 2, PTOV1, ACID2

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PTOV1 Polyclonal Antibody - Protein Information

Name PTOV1

Synonyms ACID2

Function

May activate transcription. Required for nuclear translocation of FLOT1. Promotes cell proliferation.

Cellular Location

Cytoplasm. Nucleus. Cell membrane. Cytoplasm, perinuclear region. Note=Translocates from the cytoplasm to the nucleus at the onset of S-phase (PubMed:12598323). Also localizes to lipid rafts (PubMed:15713644).

Tissue Location


Expressed in brain, heart, kidney, liver, placenta, skeletal muscle and small intestine.

PTOV1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PTOV1 Polyclonal Antibody - Images



Sample:


Liver (Mouse) Lysate at 40 ug

Primary: Anti- PTOV1 (bs-10171R) at 1/1000 dilution


Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 47 kD


Observed band size: 48 kD



Paraformaldehyde-fixed, paraffin embedded (human colon carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PTOV1) Polyclonal Antibody, Unconjugated (bs-10171R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PTOV1) Polyclonal Antibody, Unconjugated (bs-10171R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PTOV1) Polyclonal Antibody, Unconjugated (bs-10171R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.