

TRPV2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54243**Specification**

TRPV2 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	O9Y5S1
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	85981

TRPV2 Polyclonal Antibody - Additional Information**Gene ID** 51393**Other Names**

Transient receptor potential cation channel subfamily V member 2, TrpV2, Osm-9-like TRP channel 2, OTRPC2, Vanilloid receptor-like protein 1, VRL-1, TRPV2, VRL

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

TRPV2 Polyclonal Antibody - Protein Information**Name** TRPV2**Synonyms** VRL**Function**

Calcium-permeable, non-selective cation channel with an outward rectification. Seems to be regulated, at least in part, by IGF- I, PDGF and neuropeptide head activator. May transduce physical stimuli in mast cells. Activated by temperatures higher than 52 degrees Celsius; is not activated by vanilloids and acidic pH.

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:Q9WTR1}; Multi-pass membrane protein. Cytoplasm {ECO:0000250|UniProtKB:Q9WTR1}. Melanosome. Note=Translocates from the cytoplasm to the plasma membrane upon ligand stimulation (By similarity). Identified by mass spectrometry in melanosome fractions from stage I to stage IV {ECO:0000250|UniProtKB:Q9WTR1}

TRPV2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TRPV2 Polyclonal Antibody - Images