

FAM98A Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54340**Specification**

FAM98A Polyclonal Antibody - Product Information

Application	IHC-P, FC
Primary Accession	O8NCA5
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	55273

FAM98A Polyclonal Antibody - Additional Information**Gene ID** 25940**Other Names**

Protein FAM98A, FAM98A ([HGNC:24520](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=24520))

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

FAM98A Polyclonal Antibody - Protein Information**Name** FAM98A ([HGNC:24520](#))**Function**

Positively stimulates PRMT1-induced protein arginine methylation (PubMed:[28040436](http://www.uniprot.org/citations/28040436)). Involved in skeletal homeostasis (By similarity). Positively regulates lysosome peripheral distribution and ruffled border formation in osteoclasts (By similarity).

Tissue Location


Expressed strongly in colorectal cancer cells (PubMed:28040436). Expressed strongly in colorectal cancer tissues compared to wild-type colon samples (at protein level) (PubMed:28040436). Expressed strongly in colorectal cancer tissues compared to wild-type colon samples (PubMed:28040436)

FAM98A Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


FAM98A Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FAM98A) Polyclonal Antibody, Unconjugated (bs-11012R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Human breast carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FAM98A) Polyclonal Antibody, Unconjugated (bs-11012R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Blank control: hela(blue), the cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with ice-cold 90% methanol for 30 min on ice..

Isotype Control Antibody: Rabbit IgG(orange) ; Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA ; Primary Antibody Dilution: 1 μ g in 100 μ L1X PBS containing 0.5% BSA(green).