

Nicalin Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54561

Specification

Nicalin Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC
Primary Accession	O969V3
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	62974

Nicalin Polyclonal Antibody - Additional Information

Gene ID 56926

Other Names

Nicalin, Nicastrin-like protein, NCLN

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Nicalin Polyclonal Antibody - Protein Information

Name NCLN {ECO:0000303|PubMed:36261522, ECO:0000312|HGNC:HGNC:26923}

Function

Component of the multi-pass translocon (MPT) complex that mediates insertion of multi-pass membrane proteins into the lipid bilayer of membranes (PubMed:32820719, PubMed:36261522). The MPT complex takes over after the SEC61 complex: following membrane insertion of the first few transmembrane segments of proteins by the SEC61 complex, the MPT complex occludes the lateral gate of the SEC61 complex to promote insertion of subsequent transmembrane regions (PubMed:36261522). May antagonize Nodal signaling and subsequent organization of axial structures during mesodermal patterning, via its interaction with NOMO (By similarity).

Cellular Location

Endoplasmic reticulum membrane; Single-pass membrane protein

Tissue Location

Highly expressed in pancreas and skeletal muscle and, at lower levels, in heart.

Nicalin Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Nicalin Polyclonal Antibody - Images