

**Nicalin Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP54561**

## Specification

---

### Nicalin Polyclonal Antibody - Product Information

|                   |                              |
|-------------------|------------------------------|
| Application       | WB, IHC-P, IHC-F, IF, ICC, E |
| Primary Accession | <a href="#">O969V3</a>       |
| Reactivity        | Rat                          |
| Host              | Rabbit                       |
| Clonality         | Polyclonal                   |
| Calculated MW     | 62974                        |

### Nicalin Polyclonal Antibody - Additional Information

**Gene ID** 56926

#### Other Names

Nicalin, Nicastrin-like protein, NCLN

#### Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

#### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

### Nicalin Polyclonal Antibody - Protein Information

**Name** NCLN {ECO:0000303|PubMed:36261522, ECO:0000312|HGNC:HGNC:26923}

#### Function

Component of the multi-pass translocon (MPT) complex that mediates insertion of multi-pass membrane proteins into the lipid bilayer of membranes (PubMed:<a href="http://www.uniprot.org/citations/32820719" target="\_blank">32820719</a>, PubMed:<a href="http://www.uniprot.org/citations/36261522" target="\_blank">36261522</a>). The MPT complex takes over after the SEC61 complex: following membrane insertion of the first few transmembrane segments of proteins by the SEC61 complex, the MPT complex occludes the lateral gate of the SEC61 complex to promote insertion of subsequent transmembrane regions (PubMed:<a href="http://www.uniprot.org/citations/36261522" target="\_blank">36261522</a>). May antagonize Nodal signaling and subsequent organization of axial structures during mesodermal patterning, via its interaction with NOMO (By similarity).

#### Cellular Location

Endoplasmic reticulum membrane; Single-pass membrane protein

#### Tissue Location

Highly expressed in pancreas and skeletal muscle and, at lower levels, in heart.

## **Nicalin Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **Nicalin Polyclonal Antibody - Images**