

PROSAAS Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54595**Specification****PROSAAS Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q9UHG2
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27372

PROSAAS Polyclonal Antibody - Additional Information**Gene ID** 27344**Other Names**

ProSAAS, Proprotein convertase subtilisin/kexin type 1 inhibitor, Proprotein convertase 1 inhibitor, pro-SAAS, KEP, Big SAAS, b-SAAS, Little SAAS, I-SAAS, N-proSAAS, Big PEN-LEN, b-PEN-LEN, SAAS CT(1-49), PEN, Little LEN, I-LEN, Big LEN, b-LEN, SAAS CT(25-40), PCSK1N

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PROSAAS Polyclonal Antibody - Protein Information**Name** PCSK1N**Function**

May function in the control of the neuroendocrine secretory pathway. Proposed be a specific endogenous inhibitor of PCSK1. ProSAAS and Big PEN-LEN, both containing the C-terminal inhibitory domain, but not the further processed peptides reduce PCSK1 activity in the endoplasmic reticulum and Golgi. It reduces the activity of the 84 kDa form but not the autocatalytically derived 66 kDa form of PCSK1. Subsequent processing of proSAAS may eliminate the inhibition. Slows down convertase-mediated processing of proopiomelanocortin and proenkephalin. May control the intracellular timing of PCSK1 rather than its total level of activity (By similarity).

Cellular Location

Secreted {ECO:0000250|UniProtKB:Q9QXV0}. Golgi apparatus, trans-Golgi network {ECO:0000250|UniProtKB:Q9QXV0}. Note=A N-terminal processed peptide, probably Big SAAS or Little SAAS, is accumulated in cytoplasmic protein tau deposits in frontotemporal dementia and parkinsonism linked to chromosome 17 (Pick disease), Alzheimer disease and amyotrophic lateral sclerosis- parkinsonism/dementia complex 1 (Guam disease)

Tissue Location

Expressed in brain and pancreas.

PROSAAS Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PROSAAS Polyclonal Antibody - Images