

PKIB Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54937

Specification

PKIB Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O9C010
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	8468

PKIB Polyclonal Antibody - Additional Information

Gene ID 5570

Other Names

cAMP-dependent protein kinase inhibitor beta, PKI-beta, PKIB, PRKACN2

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PKIB Polyclonal Antibody - Protein Information

Name PKIB

Synonyms PRKACN2

Function

Extremely potent competitive inhibitor of cAMP-dependent protein kinase activity, this protein interacts with the catalytic subunit of the enzyme after the cAMP-induced dissociation of its regulatory chains.


PKIB Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PKIB Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PKIB) Polyclonal Antibody, Unconjugated (bs-12692R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.