

**STK32C Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP54996**

## Specification

---

### STK32C Polyclonal Antibody - Product Information

Application	IHC-P, WB
Primary Accession	<a href="#">Q86UX6</a>
Reactivity	Rat, Pig, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	54994

### STK32C Polyclonal Antibody - Additional Information

**Gene ID** 282974

#### Other Names

Serine/threonine-protein kinase 32C, 2.7.11.1, PKE, Yet another novel kinase 3, STK32C {ECO:0000312|EMBL:AAH45760.1}, YANK3

#### Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glycerol

#### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

### STK32C Polyclonal Antibody - Protein Information

**Name** STK32C {ECO:0000312|EMBL:AAH45760.1}


**Synonyms** YANK3

### STK32C Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## STK32C Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (STK32C) Polyclonal Antibody, Unconjugated (bs-12833R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



### Sample:

Molt-4(Human) Cell Lysate at 40 ug


A549(Human) Cell Lysate at 40 ug

Primary: Anti-STK32C (bs-12833R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 56 kD

Observed band size: 56 kD



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (STK32C) Polyclonal Antibody, Unconjugated (bs-12833R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.