

DNAJC9 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55041

Specification

DNAJC9 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	O8WXX5
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	29910

DNAJC9 Polyclonal Antibody - Additional Information

Gene ID 23234

Other Names

Dnaj homolog subfamily C member 9, HDJC9, Dnaj protein SB73, DNAJC9

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

DNAJC9 Polyclonal Antibody - Protein Information

Name DNAJC9

Function

Acts as a dual histone chaperone and heat shock co-chaperone (PubMed:33857403). As a histone chaperone, forms a co-chaperone complex with MCM2 and histone H3-H4 heterodimers; and may thereby assist MCM2 in histone H3-H4 heterodimer recognition and facilitate the assembly of histones into nucleosomes (PubMed:33857403). May also act as a histone co-chaperone together with TONSL (PubMed:33857403). May recruit histone chaperones ASF1A, NASP and SPT2 to histone H3-H4 heterodimers (PubMed:33857403). Also plays a role as co-chaperone of the HSP70 family of molecular chaperone proteins, such as HSPA1A, HSPA1B and HSPA8 (PubMed:17182002, PubMed:33857403). As a co-chaperone, may play a role in the recruitment of HSP70-type molecular chaperone machinery to histone H3-H4 substrates, thereby maintaining the histone structural integrity (PubMed:33857403). Exhibits activity to assemble histones onto DNA in vitro

(PubMed:33857403).

Cellular Location

Nucleus. Cytoplasm. Cell membrane. Note=Predominantly nuclear. Translocates to the cytoplasm and membrane after heat shock

Tissue Location


Expressed in heart, placenta, liver, skeletal muscle, kidney, pancreas, thymus, ovary, colon and peripheral blood

DNAJC9 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DNAJC9 Polyclonal Antibody - Images



Sample:


Thymus (Mouse) Lysate at 40 ug

Primary: Anti- DNAJC9 (bs-13024R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 30 kD

Observed band size: 35 kD



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (DNAJC9) Polyclonal Antibody, Unconjugated (bs-13024R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.