

**GARNL1 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP55122**

**Specification**

---

**GARNL1 Polyclonal Antibody - Product Information**

Application	IHC-P
Primary Accession	<a href="#">O6GYQ0</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	229832

**GARNL1 Polyclonal Antibody - Additional Information**

**Gene ID** 253959

**Other Names**

Ral GTPase-activating protein subunit alpha-1, GAP-related-interacting partner to E12, GRIPE, GTPase-activating Rap/Ran-GAP domain-like 1, Tuberin-like protein 1, p240, RALGAPA1, GARNL1, KIAA0884, TULIP1

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**GARNL1 Polyclonal Antibody - Protein Information**

**Name** RALGAPA1

**Synonyms** GARNL1, KIAA0884, TULIP1

**Function**

Catalytic subunit of the heterodimeric RalGAP1 complex which acts as a GTPase activator for the Ras-like small GTPases RALA and RALB.

**Cellular Location**

Cytoplasm. Nucleus. Note=Translocated to the nucleus, when associated with TCF3/E12

**Tissue Location**


Widely expressed..

**GARNL1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **GARNL1 Polyclonal Antibody - Images**



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GARNL1) Polyclonal Antibody, Unconjugated (bs-13286R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.