

GLI4 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55149**Specification**

GLI4 Polyclonal Antibody - Product Information

Application	IHC-P, WB
Primary Accession	P10075
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41145

GLI4 Polyclonal Antibody - Additional Information**Gene ID** 2738**Other Names**

Zinc finger protein GLI4, Krueppel-related zinc finger protein 4, Protein HKR4, GLI4, HKR4

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

GLI4 Polyclonal Antibody - Protein Information**Name** GLI4**Synonyms** HKR4**Cellular Location**

Nucleus.


GLI4 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

GLI4 Polyclonal Antibody - Images



Tissue/cell: human cervical cancer; 4% Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-GLI4 Polyclonal Antibody, Unconjugated(bs-13373R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Sample:
A549 Cell (Human) Lysate at 30 ug
Primary: Anti-GLI4 (bs-13373R) at 1/300 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 41 kD
Observed band size: 41 kD