

GOLGA7 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55177**Specification**

GOLGA7 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	O7Z5G4
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	15824

GOLGA7 Polyclonal Antibody - Additional Information**Gene ID** 51125**Other Names**

Golgin subfamily A member 7, Golgi complex-associated protein of 16 kDa, GOLGA7, GCP16

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

GOLGA7 Polyclonal Antibody - Protein Information**Name** GOLGA7**Synonyms** GCP16**Function**

May be involved in protein transport from Golgi to cell surface. The ZDHHC9-GOLGA7 complex is a palmitoyltransferase specific for HRAS and NRAS.

Cellular Location

Golgi apparatus membrane; Lipid- anchor

Tissue Location


Expressed in all tissues except colon and thymus.

GOLGA7 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

GOLGA7 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GOLGA7) Polyclonal Antibody, Unconjugated (bs-13484R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.