

ZBED3 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55196**Specification**

ZBED3 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC
Primary Accession	Q96IU2
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	25132

ZBED3 Polyclonal Antibody - Additional Information**Gene ID** 84327**Other Names**

Zinc finger BED domain-containing protein 3, Axin-interacting protein, ZBED3

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ZBED3 Polyclonal Antibody - Protein Information**Name** ZBED3**Function**

Acts as a positive regulator in the activation of the canonical Wnt/beta-catenin signaling pathway by stabilizing cytoplasmic beta-catenin (By similarity). Involved in transcription activation of Wnt target gene expression (By similarity). Plays a role in symmetric division of blastomeres in the early stages of embryogenesis via regulation of mitotic spindle central positioning and organization of the F-actin filament network (By similarity). Plays a role in regulating the distribution of cellular organelles, via modulation of cytoskeletal dynamics and cytoplasmic lattice formation (By similarity).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q9D0L1}. Membrane {ECO:0000250|UniProtKB:Q9D0L1}. Secreted

Tissue Location

Secreted in blood plasma, and expressed in skeletal muscle and adipose tissue (at protein level)

ZBED3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZBED3 Polyclonal Antibody - Images