

PCDHB10 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP55256

Specification

PCDHB10 Polyclonal Antibody - Product Information

Application	IHC-P, WB
Primary Accession	O9UN67
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	87621

PCDHB10 Polyclonal Antibody - Additional Information

Gene ID 56126

Other Names

Protocadherin beta-10, PCDH-beta-10, PCDHB10

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PCDHB10 Polyclonal Antibody - Protein Information

Name PCDHB10

Function

Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.

Cellular Location

Cell membrane; Single-pass type I membrane protein


PCDHB10 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


PCDHB10 Polyclonal Antibody - Images




Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-PCDHB10 Polyclonal Antibody, Unconjugated(bs-13725R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: human lung carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;
Incubation: Anti-PCDHB10 Polyclonal Antibody, Unconjugated(bs-13725R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (human cervical carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PCDHB10) Polyclonal Antibody, Unconjugated (bs-13725R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Sample:


A549(Human) Cell Lysate at 30 ug

Primary: Anti-PCDHB10 (bs-13725R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 84 kD

Observed band size: 112 kD

**Sample:**

HeLa(Human) Cell Lysate at 30 ug

Primary: Anti-PCDHB10 (bs-13725R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 84 kD

Observed band size: 112 kD